

The Disposal of Biosolids and Water Treatment Residuals on Soils of Arid Regions: Effect on Wheat Biomass and P and Al Availability

E.A.ELKHATIB¹ AND A.M.MAHDY¹

1-Department of Soil and Water, College of Agriculture,
Alexandria University,
Alexandria-Elshatby, 21545
EGYPT

amahdy73@yahoo.com;amahdy73@gmail.com;selkhatib1@yahoo.com

ABSTRACT:-Land co-application of biosolids and WTR is a new concept. Therefore, information on the effect of co-application of biosolids and WTR on plant growth and elements uptake are very limited especially in alkaline soils. A glasshouse experiments was established to evaluate the effects of co-application of WTR and biosolids on agronomic performance of wheat crop grown in alkaline soils as well as P and Al plant concentration and uptake, and to improve management of industrial and toxic wastes and provides environmentally sound guidelines for their disposal. The results indicated that increases of 47, 359 and 55 % in total dry matter yield were achieved as a result of applying 40 gkg⁻¹ WTR and 10 gkg⁻¹ biosolids to clay, sandy and calcareous soils respectively. In all studied soils treated with a constant biosolid rate 10 gkg⁻¹, application of 20 gkg⁻¹ WTR significantly increased plant P concentration in the plant materials. Combined analyses of all soils ,all treatments of biosolid and WTR rates studied indicated clearly significant relationships between ABDTPA P concentration and P uptake ($r = 0.81$, $p < 0.001$). Application of WTR combined with biosolid showed significant increases in extractable Al in all the soils studied. However, application rates of WTR and biosolid did not increase extractable Al in amended soils beyond 6.06 mg Al kg⁻¹.

Keywords: - Biosolids, Water treatment residuals, Phosphorus, Aluminum, Availability

1 Introduction

Alum [Al₂ (SO₄)₃. 14 H₂O] is commonly used in the municipal water treatment process to destabilize colloids for subsequent flocculation and water clarification. Water treatment residuals (WTR) are by-products of water purification systems in which undesirable attributes of the raw water such as turbidity, color and dissolved solids are removed by a variety of physical and chemical processes. WTR have commonly either been returned to the source waterway downstream of the treatment plant intake or has been released to a municipal sanitary sewer. Both methods of disposal of water treatment residuals (WTR) have become unattractive for a variety of reasons. As the alternatives to disposal of WTR have decreased in recent years, more attention has focused on beneficial reuse of the material. One such beneficial reuse is land application. However, the potential benefits of applying WTR to the soil have been limited due to its postulated reduction of plant available P and potential plant Al toxicity with increasing WTR rates. Co-application of WTR with biosolids inherently high in P could offer a good opportunity to reconcile these problems. Land co-application of biosolids and WTR is a new concept. Therefore, information on the effect of co-application of biosolids and WTR on plant growth and elements uptake are very

limited especially in alkaline soils. Co-application of WTR with biosolids may be advantageous in terms of a cost saving and potential reduction of bioavailable P in high P containing sludge [1].

2 Problem Formulation

Agricultural land application of biosolid waste materials is known to be an effective means of recycling organic matter and plant nutrients from waste materials. Biosolids are a valuable source of plant nutrients and organic matter, but it also contains very high levels of P and various levels of heavy metals. The excessive P could cause undesirable environmental impacts, and threaten surface or ground water qualities with eutrophication-associated problems. Land application of biosolids may also significantly increase heavy metals accumulation in agricultural soils [2].

Water treatment plants have thousands of tons of alum-laden lime sludge available for land application which are mostly disposed in landfills. As landfills space becomes less available, it is urgent to investigate alternative approaches of disposing of water treatment residuals(WTR).Concern over agricultural land application of WTR are due to its reduction of plant

available P and potential Al toxicity with increasing WTR rates [3]. However, coapplication of WTR and P-laden biosolids may offer an opportunity to effectively reconcile these environmental problems. The combination of WTR and biosolids promotes agricultural land application of non-hazardous wastes and recycle available organic matter and plant nutrients that accumulate in biosolids.

3 Problem Solution

To study and solve the problem, the objectives of this study were: to evaluate the effects of co-application of WTR and biosolids on agronomic performance of wheat crop grown in alkaline soils as well as P plant concentration and uptake and to improve management of industrial and toxic wastes and provides environmentally sound guidelines for their disposal. To achieve these objectives, incubation and glasshouse experiments were conducted as follows: Three soils with different properties (clay, sandy and calcareous) were selected for the study and sampled (0-15 cm) from three different locations. Sub-samples of the air-dried soils, WTR and biosolids were ground to pass a 2-mm sieve prior to the following chemical analysis: Soil pH and EC were measured in the soil-paste extracts, but pH and EC of WTR and biosolids were measured in 1: 2 sludge/deionized water [4]. The organic matter content was determined by dichromate oxidation [5], cation exchange capacity was determined by IM NaOAc [6]. Particle size analysis was determined by the hydrometer [7]. Calcium carbonate content was determined using a calcimeter [8]. Total nitrogen was determined by the Kjeldahl/digestion method [9]. Available P was extracted by AB-DTPA test [10]. Total Al was determined using the acid ammonium oxalate method [11]. Extractable Aluminum was extracted by 1M KCl [12] and determined colorimetrically by 8-hydroxyquinoline butyl acetate method [13]. The selected properties of the three soils are summarized in Table 1.

To ensure amendment-soils equilibria, incubation experiment was conducted. Four biosolids rates (0, 10, 20 and 30 g kg⁻¹ on an oven dry basis) and/or five WTR rates (0, 10, 20, 30 and 40 g kg⁻¹) and/or co-application rates of WTR and rates of biosolids were applied to each soil (calcareous, sandy and clay soils) thoroughly mixed and placed in Jars (2 kg). Following amendments applications, the soil water content was brought to field capacity. Jars were covered with perforated plastic cover and incubated at 25°C for 60 days.

Seeds of wheat (*Triticum aestivum*) cultivar were sown in pots containing 2 kg of soil (s) with co-application rates of WTR and biosolids. The seedlings were thinned to 4 seedlings per pot and distilled water

was added to bring the soil moisture to 70% of field capacity. The experiment was arranged in split-split plot design with four replicates. Plants were harvested after 13 weeks. Plant shoots, panicles and roots were harvested separately, oven dried at 65°C for 48 h to determine dry matter yield. Subsamples of ground plant material were ashed in muffle furnace at 450°C for 6h, and analyzed colorimetrically for P [14]. The ammonium bicarbonate-DTPA extractant solution was used to extract available phosphorus from soils treated with and without WTR and biosolids after harvest [8]. Statistical and mathematical analyses were performed using Statistical Analysis System [15]. Analysis of Variance (ANOVA) techniques was used to determine treatment effects and check for interaction. The least significant difference method was used to separate treatment means. Regression analysis was employed to determine the relationships between available P concentration in soils and P concentration in plants. The results of current study were:

3.1 Soils, (WTR) and biosolids characteristics

Selected properties of the soils, WTR and biosolids used in the study are given in Table (1). The soils differ dramatically in their textures, CaCO₃ and organic matter contents. The sandy soil samples represent soil with coarse texture, low contents of CaCO₃ and organic matter (O.M). It is classified as (*Typic Torripsamments*). In contrast, the clay soil is (*Typic Torrifluvents*), containing approximately 3 to 10 times as much as clay and organic matter contents. The CaCO₃ content and the CEC are much higher than the sandy soil. The pH of the clay soil is 0.5 unit higher than the sandy soil. The calcareous soil is classified as (*Typic Calciorthids*). The calcium carbonate content in the calcareous soil samples is 6 times higher than that in the clay soil samples. The three studied soils had concentrations of ABDTPA-P ranging from low (sandy soil) to high (clay soil). The clay soil contains approximately 2.5 and 1.5 times ABDTPA-P concentration more than that of the sandy and calcareous soils, respectively.

The WTR was slightly alkaline (7.45) within the adequate typical range for plant growth (5-8) [16]. The EC of WTR is well below the 4 dSm⁻¹ associated with the high exchange capacity of the WTR indicates its ability to supply cationic nutrients for plant growth. The organic matter content of the WTR is considerably greater than typical levels in soils of arid ecosystems. The small amount water soluble P (< 0.04 % of the total P) extracted from WTR implied strong P binding by the WTR. Dayton et al. [17] reported that low P extractability of WTR was due to the abundance of Al. However, the ABDTPA-P concentration in WTR was very similar.

Table1. Some physical and chemical characteristics of studied soils

Characteristics	Units	Clay	Sandy	Calcareous	WTR	Biosolids
EC*	dSm ⁻¹	2.66	3.84	2.92	1.67	11.25
PH*		8.13	7.69	8.08	7.45	6.69
CaCO ₃	%	5.79	0.24	35.68	-	-
Sand	%	59.64	86.82	74.00	-	-
Silt	%	14.13	2.51	10.15	-	-
Clay	%	26.23	10.67	15.85	-	-
Texture		S.C.L	L.S	S.L	-	-
O.M	%	0.85	0.10	0.46	5.70	45.00
T-N	%	0.22	0.03	0.09	0.42	3.20
T-P	%	0.09	0.03	0.05	0.19	0.46
T-Al	g kg ⁻¹				38.01	3.14
CEC	Cmol(+)kg ⁻¹	39.13	8.70	26.00	34.78	73.57
AB-DTPA-P	mg kg ⁻¹	8.13	3.12	5.15	8.32	24.00
Extractable Al	mg kg ⁻¹	1.03	0.13	0.08	28.18	4.22
Soil solution-P	mg kg ⁻¹	1.98	0.89	1.22	0.73	2.13
Soil solution-Al	mg kg ⁻¹	0.03	0.01	0.02	1.80	0.18
W.H.C	g kg ⁻¹	259.30	93.80	166.70	470.00	250.00

SCI:Sandy Clay Loam; LS:Loamy Sand; SL:Sandy Loam

The water holding capacity of WTR is high (470 gkg⁻¹). Therefore, the WTR could be considered a good ameliorating agent to soil properties [18].

Biosolid is slightly acidic with high content of organic matter. Biosolids could be regarded as a low analysis P source (0.46 %) but the AB-DTPA extractability suggests that total P may not completely assess P solubility. The total nitrogen and phosphorus was higher than WTR. The water holding capacity of biosolid (250 gkg⁻¹) was lower than its value in WTR. Therefore, the coapplication of WTR and biosolids could be considered good ameliorating agents to soil properties.

3.2 Dry matter yield of wheat

The effect of WTR rates, co-applied with different biosolids rates on total dry matter of wheat grown on the three studied soils is shown in Table (2). In the all studied soils treated with WTR rates co-applied with 10 gkg⁻¹ biosolids rate, the total dry matter yield was not significantly different between the control treatment and the 10 gkg⁻¹ WTR treatment. However, a significant increase in total dry matter was found between the control treatment and 20 or 30 or 40 gkg⁻¹ treatments. Increases of 47, 359 and 55 % in total dry matter yield were achieved as a result of applying 40 gkg⁻¹ WTR and 10 gkg⁻¹ biosolids to clay, sandy and calcareous soils respectively. However, increases of 52, 343, and 58 % in total dry matter yield were achieved as a result of applying 30 gkg⁻¹ WTR and 20 gkg⁻¹ biosolids to clay, sandy, and calcareous respectively. Increases of 55, 403 and 72 % in total dry matter yield were achieved as a result of applying 30 gkg⁻¹ WTR and 30 gkg⁻¹ biosolids

to clay, sandy and calcareous soils respectively. In the soils treated with WTR rates co-applied with 20 gkg⁻¹ biosolid rate, the total dry matter yield was significantly different between the control treatment and the 20,30 or 40 gkg⁻¹ treatments. These results coincide with the results of Harris-Pierce et al., [19,20]. Heil and Barbarick [21] also observed an increase in dry matter with WTR application at high rates. In sandy and calcareous soils treated with WTR coapplied with a constant biosolid rate of 30 gkg⁻¹, there was no significant different between the control treatment and that the 10 gkg⁻¹ treatment. However, there was a significant different between the control treatment and 20 or 30 gkg⁻¹ treatments. Soil, biosolid treatments and WTR rates main effects were significant for total dry yield ($p < 0.001$) (Table 2). According to the previous results, it can be concluded that the application of WTR to high P soils may be a very good opportunity for farmers and water municipalities to reconcile several problems. Many farmers are being pressured to reduce the pollution impact of their traditional fertilization practices are less threatened. Additionally, farmers would receive a very good conditioner material to improve the physical and chemical characteristics of soils and, consequently high production of yield. Water utilities could have a more economic and labor conservative disposal method than the more common methods of WTR disposal, such as land filling, sewage disposal and coagulant recovery. Finally, the co-application of WTR and biosolids, resulted in total dry matter yield more than WTR or biosolids when it is applied individually [1].

Table 2. Total dry matter yield of wheat plants grown in the three soils as influenced by co- application of biosolids and WTR rates

Biosolids rate, gkg ⁻¹	WTR rate, gkg ⁻¹	Total dry matter yield, g pot ⁻¹		
		Clay	Sand	Calcareous
10	0	2.71	2.54	2.83
10	10	2.91	2.80	2.87
10	20	3.24	3.00	3.10
10	30	3.34	3.14	3.22
10	40	3.71	3.49	3.49
	LSD_{0.05}	0.42	0.41	0.45
20	0	3.11	2.83	3.03
20	10	3.31	2.74	2.92
20	20	3.56	3.15	3.29
20	30	3.85	3.37	3.55
20	40	3.37	3.23	3.28
	LSD_{0.05}	0.30	0.28	0.64
30	0	3.40	3.08	3.22
30	10	3.49	3.26	3.38
30	20	3.70	3.53	3.59
30	30	3.92	3.82	3.88
30	40	3.43	3.21	3.37
	LSD_{0.05}	0.64	0.39	0.30
Analysis of variance		F-test		
Soil (S)		*		
Treatment (T)		***		
Rate (R)		***		
T X S		NS		
R X S		NS		
R X T		***		
R X T X S		NS		

*,*** significant at the 0.05 and 0.001 probability levels respectively.

NS: Not Significant.

3.3 Phosphorus concentration and uptake

In general, phosphorus concentration in plants tends to be accumulated in the order panicles > shoots > roots (Table 3). Significant soil type, biosolid treatments, WTR rates and their interactions effects were found for phosphorus concentration in panicles, shoots and roots of wheat plants grown in all the soils studied (Table 3). In all studied soils treated with a constant biosolid rate 10 gkg⁻¹, application of 20 gkg⁻¹ WTR significantly increased plant P concentration in the plant materials (Table 3). The P concentration significantly increased in plant materials (e.g.panicles) from 3000 to 3504, from 2910 to 3180 and from 2902 to 3030 mgkg⁻¹ in clay, sandy and calcareous soils treated with 20 gkg⁻¹ biosolid and WTR rates (Table 3). The increase in extractable P in the amended soil with increasing the application rate of WTRs might be due to the high content of available P in the WTRs used in this study. Therefore the addition of cations from the WTR-application (10-30 g.kg⁻¹) was not able to effectively reduce the extractable P in the soils, which might be attributed partially to the inaccessibility to P held on intraparticle sites [22] and also more reaction time might be required to reach the equilibrium between the cation of WTRs and extractable P of biosolids within the treated soils [22]. However, further increase in WTR application rate has resulted in negative significant impact on plant P concentration. The application of the highest sludge rate (30 gkg⁻¹) produced higher P concentration in plant materials than the other two lower treatments. These results coincide with the results of Heil and Barbarick [21] who indicated that WTR have a high capacity to fix P and that plant P deficiencies develop

when plants are grown in WTR-soil mixtures or coapplied with a constant rate of biosolids.

Similar to wheat phosphorus content data, P uptake increased at the low WTR rates coapplied with 10 or 20 or 30 gkg⁻¹ biosolid treatments (Table 3). Soil type, biosolid treatment, WTR rates and their interactions significantly affected P uptake. The P uptake was higher in clay soils than in sandy and calcareous soils at all biosolids treatments coapplied with WTR rates, but the 20 gkg⁻¹ WTR coapplied with 30 gkg⁻¹ biosolid treatment was the best co-application rate (2:3 ratio). Such data indicated that co-mixing of WTR and biosolids at ratios of 4:1 will adsorb all soluble biosolids P, and beyond this ratio the WTR could adsorb all biosolids available P and possibly some soil-borne P [1].

3.4 Phosphorus extractability after wheat harvest

Soil type, biosolid treatments, WTR rates and their interactions significantly affected AB-DTPA extractable P (Table 4). Application of WTR at rates of 10, 20 and 30 g kg⁻¹ to clay and sandy soils treated with 10 gkg⁻¹ biosolid, significantly increased AB-DTPA extractable P. In calcareous soil the extractable P increased with increasing WTR up to 20 gkg⁻¹ and decreased with increasing WTR application rate. In all the soils studied treated with WTR co-applied with 20 or 30 gkg⁻¹ biosolid, the extractable P significantly increased at rates 10 and 20 gkg⁻¹ WTR, then the extractable P dramatically decreased to about 35% compared with the control treatment (Table 4). The use of WTR as a soil or poultry litter amendment have been reported to significantly lower extractable P concentrations [22].

Table3. Phosphorus concentrations and uptake of wheat plants grown in the three soils as affected by co-application of biosolids and WTR rates

Biosolids rate	WTR rate	Clay				Sandy				Calcareous			
		gkg ⁻¹	PP	SP	RP	PU	PP	SP	RP	PU	PP	SP	RP
			mg.kg ⁻¹			mg.pot ⁻¹				mg.kg ⁻¹			mg.pot ⁻¹
10	0	2889.00	2003.00	1713.00	6.82	2801.00	1998.00	1415.00	6.27	2714.00	2101.00	1571.00	6.86
10	10	3020.00	1801.00	1780.00	7.26	2810.00	2000.00	1530.00	6.85	2736.00	2200.00	1600.00	7.10
10	20	3309.00	1906.00	1830.00	8.83	2990.00	2100.00	1770.00	7.77	2813.00	2391.00	1690.00	8.00
10	30	2811.00	1330.00	1260.00	7.02	2920.00	1550.00	1410.00	7.16	2501.00	2010.00	1201.00	7.12
10	40	2213.00	960.00	910.00	6.06	2660.00	630.00	510.00	5.99	2101.00	1830.00	410.00	6.67
	LSD _{0.05}	14.37	9.46	15.43	1.22	15.37	22.64	18.98	0.90	18.83	23.76	13.59	1.03
20	0	3000.00	2643.00	2180.00	8.81	2910.00	2315.00	2214.00	7.55	2902.00	2440.00	1857.00	8.09
20	10	3109.00	2020.00	1801.00	8.79	3020.00	2081.00	1606.00	7.03	2881.00	1880.00	1670.00	7.09
20	20	3504.00	2504.00	1880.00	11.03	3180.00	2220.00	1690.00	8.38	3030.00	1990.00	1717.00	8.27
20	30	2603.00	1440.00	1202.00	8.01	2690.00	1370.00	930.00	6.72	2201.00	1101.00	810.00	5.87
20	40	2091.00	710.00	580.00	5.14	2310.00	580.00	440.00	4.61	1681.00	630.00	507.00	3.89
	LSD _{0.05}	31.61	15.53	17.64	0.76	31.38	24.90	16.65	0.60	19.07	27.88	22.99	1.54
30	0	4786.00	2990.00	2500.00	13.53	3501.00	2612.00	2402.00	9.60	3334.00	2701.00	2423.00	9.83
30	10	4880.00	2201.00	1890.00	12.94	3440.00	2103.00	1630.00	9.16	2902.00	1803.00	1499.00	8.12
30	20	5010.00	2403.00	1901.00	14.25	3630.00	2221.00	1690.00	10.51	3011.00	1880.00	1560.00	8.95
30	30	2710.00	1101.00	710.00	7.63	2803.00	990.00	801.00	7.52	1801.00	621.00	570.00	4.89
30	40	1991.00	603.00	499.00	4.64	2002.23	510.00	460.00	4.33	1590.00	460.00	420.00	3.63
	LSD _{0.05}	33.19	24.36	81.36	3.05	52.78	26.66	24.09	0.80	35.79	21.65	21.81	0.53
Analysis of variance			PP			SP				RP			PU
Soil (S)			***			***				***			***
Treatment (T)			***			***				***			***
Rate (R)			***			***				***			***
T X S			***			***				***			***
R X S			***			***				***			***
R X T			***			***				***			***
R X T X S			***			***				***			*

*** significant at the 0.05 and 0.001 probability levels respectively.
 PP: panicles phosphorus SP: shoots phosphorus
 RP: root phosphorus PU: phosphorus uptake

Table 4. AB-DTPA extractable phosphorus concentrations for three soils influenced by co-application of biosolid and WTR rates

Biosolids rate	WTR rate	AB-DTPA extractable P , mg.kg ⁻¹			
		gkg ⁻¹	Clay	Sandy	Calcareous
10	0		6.87	8.13	8.27
10	10		10.13	8.88	9.23
10	20		12.19	9.12	11.77
10	30		13.99	11.14	7.13
10	40		7.66	6.12	4.22
	LSD _{0.05}		0.82	0.82	1.18
20	0		12.66	12.42	13.01
20	10		14.28	12.93	13.86
20	20		19.09	15.87	15.08
20	30		12.22	8.18	9.01
20	40		5.88	4.43	5.82
	LSD _{0.05}		1.03	0.88	0.43
30	0		17.99	13.99	15.83
30	10		22.18	18.82	19.39
30	20		26.27	21.23	23.13
30	30		15.55	11.98	12.91
30	40		6.22	5.82	6.02
	LSD _{0.05}		0.53	0.70	0.53
Analysis of variance				F-test	
Soil (S)				AB-DTPA extractable P	***
Treatment (T)					***
Rate (R)					***
T X S					***
R X S					***
R X T					***
R X T X S					***

*** Significant at the 0.001 probability level.

Codling et al.[23], Elliott et al. [24], and other researchers noted similar declines in soil P concentration after the addition of WTRs to manure-treated soils. Combined analyses of all soils, all treatments of biosolid and WTR rates studied revealed clearly significant

relationships between AB-DTPA extractable P concentration and P uptake (r = 0.81, p < 0.001, fig.1). These results agree with the studies of Harris-Pierce et al. [19] and Shreve et al. [25].




Fig. 1. Relationship between AB-DTPA P and P uptake of wheat plants grown in biosolids-WTR-treated soils .

The polynomial quadratic model was used to describe the relationship between P concentration in AB-DTPA extracts and co-application of WTR and biosolid rates (Fig.2).

Fig.2: Polynomial quadratic models of extractable P in the three soils treated with WTR coapplied with constant rates of biosolid. *, ** Significant at the 0.05 and 0.01 probability levels respectively

3.5 Al concentration and uptake

Aluminum concentration tends to be accumulated in the following order: roots > shoots > panicles in all studied soils treated with biosolids and different rates of WTR (Table 5). Root staining techniques have shown that Al accumulates principally in the root tips of the main root and lateral root tissue (26).

The soil type, biosolid treatments, WTR rates and their interactions significantly affected plant Al concentration in panicles (Table 5). However, soil type, biosolid treatments and WTR rates significantly affected shoot Al and root Al concentrations (Table 5). Aluminum concentrations in panicles, shoots and roots of wheat plants grown in the three soils treated with biosolids increased in a stepwise fashion with increasing WTR (Table 5). The increased Al in soils treated with WTR coapplied with 30gkg⁻¹ biosolid was higher than Al in soils treated with 20 or 10 gkg⁻¹ biosolids. However, Al concentrations in all plant parts are well below the maximum tolerable levels of dietary Al for domestic animals (200 to 1000 mg kg⁻¹) and humans (27).

The total Al uptake was significantly affected by soil type, biosolid treatments, WTR rates and their interactions. In general, application of WTR in a rate of 10 g kg⁻¹ coapplied with 10, 20 and 30 gkg⁻¹ biosolid rates did not significantly affect Al uptake. However, addition of 30 or 40 g kg⁻¹ WTR more than doubled the total Al uptake from all soils compared to the 10 g kg⁻¹ rate (Table 5).

3.6 Aluminum extractability after wheat harvest

Extractable Al concentrations in soils as affected by different rates of WTR and biosolid application are shown in (Table 6). Soil type, biosolid treatments and WTR rate significantly affected extractable Al (Table 6). Application of WTR combined with biosolid showed significant increases in extractable Al in all the soils studied. However, application rates of WTR and biosolid did not increase extractable Al in amended soils beyond 6.06 mg Al kg⁻¹ (Table 6). Adverse effects are associated with much higher levels of extractable Al (> 60 mg Al kg⁻¹) for wheat (28) Sloan et al., 1995). Therefore slight increases in extractable Al from co-application of WTR and biosolid should not have adverse effects on soils or plants.




Table (5): Aluminum concentrations and uptake of wheat plants grown in the three soils by as affected by co-application of biosolid and WTR rates.

Biosolids Treat.	WTR rate	Clay				Sandy				Calcareous							
		PA	SA	RA	AU	PA	SA	RA	AU	PA	SA	RA	AU				
	gkg ⁻¹		mg.kg ⁻¹		mg.pot ⁻¹		mg.kg ⁻¹		mg.pot ⁻¹		mg.kg ⁻¹		mg.pot ⁻¹				
10	0	1.00	2.31	6.51	4.82	0.41	0.87	1.70	1.59	0.60	0.83	1.20	2.02				
10	10	1.09	3.80	7.60	7.19	0.79	1.20	2.71	2.93	0.72	1.03	2.49	2.65				
10	20	1.26	4.98	10.33	10.72	0.98	1.51	5.01	4.27	0.77	1.30	4.12	3.65				
10	30	1.88	5.18	16.82	14.12	2.16	5.12	11.11	12.47	0.99	4.37	9.95	9.66				
10	40	2.57	8.98	21.19	22.69	2.18	6.00	14.73	15.24	3.13	6.07	16.33	17.24				
	LSD _{0.05}	0.62	0.90	1.07	1.69	0.30	0.28	1.51	1.87	0.62	0.38	1.49	1.75				
20	0	1.09	2.33	7.78	5.85	0.48	0.92	2.18	2.03	0.66	0.89	1.99	2.44				
20	10	1.13	3.92	8.16	8.25	0.80	1.25	3.03	3.01	0.75	1.11	3.35	3.01				
20	20	1.43	5.22	11.93	12.48	1.01	1.55	5.71	4.86	0.81	1.39	4.70	4.27				
20	30	2.00	5.60	17.08	16.80	2.40	6.00	11.95	15.49	1.02	5.01	11.08	11.84				
20	40	3.03	8.99	23.12	21.27	2.43	7.12	15.53	17.17	3.70	6.99	18.41	19.50				
	LSD _{0.05}	0.36	1.01	2.26	3.14	0.25	1.32	0.84	2.49	0.22	1.29	1.67	1.76				
30	0	1.35	3.18	6.81	8.07	0.66	1.83	3.08	3.74	1.93	1.99	3.12	6.50				
30	10	1.41	4.02	9.12	10.01	0.83	2.55	3.56	5.43	1.95	2.30	3.93	7.45				
30	20	1.66	5.28	13.14	13.83	1.18	2.62	6.08	7.14	2.85	2.51	5.12	10.31				
30	30	2.40	6.09	18.19	19.55	2.81	7.17	13.14	19.89	3.18	6.04	13.08	19.36				
30	40	3.81	9.13	25.16	25.34	2.96	8.19	17.17	18.79	4.03	7.33	15.66	20.64				
	LSD _{0.05}	0.26	0.89	2.48	2.86	0.44	0.90	1.43	3.21	0.24	0.46	1.76	2.20				
F-test																	
<i>Analysis of variance</i>		PA				SA				RA				AU			
Soil (S)		***				***				***				***			
Treatment (T)		***				***				***				***			
Rate (R)		***				***				***				***			
T X S		***				NS				NS				*			
R X S		***				***				***				***			
R X T		***				NS				NS				***			
R X T X S		***				NS				**				*			

*, **, *** significant at the 0.05, 0.01 and 0.001 probability levels respectively.

NS: non significant

PA: panicles aluminum SA: shoots aluminum RA: roots aluminum AU: aluminum uptake

Table (6): Extractable and soil solution aluminum concentrations for three soils influenced by co-application of biosolid and WTR rates.

Biosolids treatment	WTR rate	Extractable Al, mg.kg ⁻¹		
		Clay	Sandy	Calcareous
	gkg ⁻¹			
10	0	1.09	1.06	0.93
10	10	1.91	1.86	1.18
10	20	2.78	2.61	2.06
10	30	4.09	4.13	2.77
10	40	5.71	5.88	3.98
	LSD _{0.05}	0.26	0.32	0.55
20	0	1.82	1.73	1.09
20	10	1.92	1.99	1.43
20	20	2.88	2.70	2.98
20	30	4.03	4.80	3.18
20	40	5.93	5.91	4.16
	LSD _{0.05}	0.20	0.31	0.23
30	0	2.22	2.71	2.11
30	10	2.37	2.08	1.57
30	20	2.98	2.87	3.13
30	30	4.74	4.88	3.45
30	40	6.06	5.99	4.90
	LSD _{0.05}	0.27	0.47	0.39
<i>Analysis of variance</i>		F-test		
		Extractable Al		
Soil (S)		***		
Treatment (T)		***		
Rate (R)		***		
T X S		NS		
R X S		***		
R X T		***		
R X T X S		***		

*** Significant at the probability level 0.001.

NS: Not Significant

Combined analyses of soils, biosolid treatment and WTR rates showed significant relationship between KCl

extractable Al and Al uptake ($r = 0.89, P < 0.001$, Fig.2).

3.7 soil pH

rates of biosolid. *, ** Significant at the 0.05 and 0.01 probability levels respectively




Fig.(2): Relationship between extractable Al and Al uptake of wheat plants grown in biosolids-WTR-treated soils.

The polynomial quadratic model was used to describe the relationship between WTR co-applied with biosolid rates and the extractable Al concentration in the three soils studied (Fig.3).

Application of WTR to clay soil showed decreases in soil solution pH from 7.23 to 7.13 at a constant rate 10 g kg⁻¹ biosolid and from 7.10 to 7.05 at a constant rate 20 g kg⁻¹ biosolid and from 7.33 to 6.94 at a constant rate 30 g kg⁻¹ biosolid (fig 4A). In general, application of WTR combined with biosolid rates significantly decreased soil solution pH of the studied soils (fig 4). These results in agreement with McBride,1994 (29)) who reported that as the pH is lowered, the concentration of soluble aluminum, which is toxic, increases. Once soil pH is lowered below 5.5 , aluminosilicate clays and aluminum hydroxide minerals begin to dissolve, releasing aluminum-hydroxy cations and Al (H₂O)₆³⁺ that then exchange with other cations from soil colloids (29).









Fig.3: Polynomial quadratic models of extractable Al in the three soils treated with WTR coapplied with constant

Fig. (5): Effect of WTR and biosolids additions on pH of the clay (A), sandy(B) and calcareous(C) soils.

4 Conclusions

The low application rates of WTR produced greater plant dry matter yield associated with the three studied soils without creating a deficiency of phosphorus. Apparently, P fixation was minimal at these rates. The low application rates of the WTR used in this study could be safely applied to the studied soils. Based on our experiment results, the ratio of 2:3 WTR/biosolids is considered the best application ratio because of its positive effects on plant dry matter. Co-application of WTR with biosolids may be advantageous in terms of a cost saving and potential reduction of bioavailable P in high P containing biosolids; however, more research and additional studies are necessary to confirm these results under field conditions.

5 Acknowledgements

The authors gratefully acknowledge the financial support provided by Consortium on Science, Technology and Innovation for the South (Formerly, TWNSO) to E.A. Elkhatib (JRP n.07/02).

References:

- [1] J.A.Ippolito, K.A. Barbarick, and E.F.Rendte , Co-application of water treatment residuals and biosolids on two range grasses. *J. Environ. Qual.* Vol.28, 1999, pp1644-1650.
- [2] J.J.Sloan,R.H.Dowdy,,M.S.Dolan,and R.Linden,.Longterm effects of biosolids application on heavy metals bioavailability in agricultural soils.*J.Environ.Qual*, Vol 26, 1997,pp966-974.
- [3] J.F. Power and W.A. Dick, *Land application of agricultural, industrial, and municipal by-products*. ASA, CSSA, and SSSA, Madison, WI,2000.
- [4] L.A .Richards,*Diagnosis and Improvement of Saline and Alkaline Soils*.USDA.Handbook 60.US Government Printing Office,Washington,D.C.1954.
- [5] D.W.Nelson,and L.E.Sommers,*Total carbon, organic carbon and organic matter*,pp.539-549.In A.L.Page, R.H.Miller, and D.R. Keeney(eds). *Methods of Soil Analysis*, American society of agronomy, Madison, Wisconsin, USA, 1982.
- [6] J.D. Rhoades, *Cation exchange capacity* In A.L. Page , R.H. Miller and D.R Keeney(eds). *Methods of soil analysis.*, American society of agronomy, Madison, Wisconsin, USA,1982.
- [7] P.R. Day, . *Particle fraction and particle size analysis* Pp:545-566.InA.C.Black,D.D.Evans, L.E.Ensminger, J.L.White, and F.E.Clark.(eds)*Methods of Soil Analysis*. Pat I. American society of agronomy, Madison, Wisconsin, USA.1965.
- [8] R.E. Nelson, *Carbonate and gypsum*.pp 181-197. In A.L. Page R.H.Miller, and D.R. Keeney(eds). *Methods of Soil Analysis*, American society of agronomy, Madison, Wisconsin, USA.1982.
- [9]J.M. Bremner and C.S.Mulvaney, Nitrogen-total.pp595 In A.L. Page R.H.Miller, and D.R. Keeney(eds). *Methods of Soil Analysis*, American society of agronomy, Madison, Wisconsin, USA.1982.
- [10] P.N.Soltanpour,and A.P. Schwab,New soils test for simultaneous of macro and micronutrients in alkaline soils, *Commun. Soil Sci. Plant Anal.*, Vol. 3, 1977,pp.195–207.
- [11]G.J. Ross, and C. Wang,*Acid ammonium oxalate method*.In M.R.Carter (ed) *Soil sampling and methods of analysis*.Lewis Pub.,Ann Arbor,MI.1993.
- [12] R. Barnhisel and P.M. Bertsch,Aluminum,pp275 . In A.L.Page, R.H.Miller, and D.R. Keeney(eds). *Methods of Soil Analysis*, American society of agronomy, Madison, Wisconsin, USA, 1982.
- [13] P.R Bloom, R.Mr Weave, and M.B. McBride, The spectrophotometric and flurometric determination of aluminum with 8-hydroxyquinoline and butyl acetateextraction.*SoilSci.Soc.Am.J*.Vol42, 1978,pp712-716.
- [14] J.B. Jones,*Laboratory guide of conducting soil tests and plant analysis*.CRC Press. New York, Washington,D.C.USA.2001

- [15] SAS Institute, *SAS/STAT User's guide*. Version 6.4th ed. SAS Inst., Cary, N.C. 1994.
- [16] H.L. Bohn, B.L. McNeal, and G.A.P. O'Connor, *Soil Chemistry* 2nd Ed. Wiley, New York. 1985
- [17] E.A. Dayton, N.T. Basta, Jakober, and J.A. Hattey, Using treatment residuals to reduce phosphorus in agricultural runoff, *J. Am. Water Works Assoc.*, Vol. 95, No. 4, 2003 pp. 151–157.
- [18] T.M. Skene, J.M. Oades, and G. Kilmore, Water treatment sludge: A potential plant growth medium. *Soil Use and Management*, Vol. 11, 1995, pp 29-33.
- [19] R. Harris-Pierce, K.A. Barbarick, and E.F. Redente. The effect of sewage sludge application on native rangeland soils and vegetation. *Annual Rep.* Meadow Springs Ranch, Fort Collins, CO. 1994
- [20] R. Harris-Pierce, K.A. Barbarick, and E.F. Redente. The effect of sewage sludge application on native rangeland soils and vegetation *Annual Rep.* Meadow Springs Ranch, Fort Collins, CO. 1993
- [26] E. Matsumoto, F. Hiasawa, H. Torikai, and E. Takakaski., Localization of absorbed aluminum in pea roots and its binding to nucleic acid. *Plant And Cell Physiol.* 17, 1976., pp 127-137.
- [27] National Research Council, Mineral tolerance of domestic animals. Natl. Academy of Sciences, Washington, D. C. 1980.
- [28] J. Sloan, N.T. Basta, and R.L. Westerman. Aluminum transformations and solution equilibria induced by banded phosphorus fertilizer in acid soil. *Soil Sci. Soc. Am. J.* 59, 1995. pp 327-364.
- [29] M.B. McBride, *Environmental chemistry of soils*. Oxford University press. New York. , 1994..
- [21] D.N. Heil and K.A. Barbarick, Water treatment sludge influence on the growth of sorghum-sudangrass. *J. Environ. Qual.*, Vol. 18, 1989, pp 292-298.
- [22] K.C. Makris, H. El-shall, W.G. Harris, G.A. O'Connor, and T.A. Obreza, Intraparticle phosphorus diffusion in a drinking water treatment residual at room temperatures. *J. Colloid Interface Sci.*, Vol. 277, 2004, pp 417-424.
- [23] E.E. Codling, R.L. Chaney, and C.L. Mulchi, Use of aluminum and iron-rich residues to immobilize phosphorus in poultry litter and litter amended soils. *J. Environ. Qual.* Vol. 29, 2000, pp 1924-1931.
- [24] H.A. Elliott, G.A. O'Connor, P. Lu, and S. Brinton, Influence of water treatment residuals on phosphorus solubility and leaching. *J. Environ. Qual.* Vol. 31, 2002, pp 1362-1369.
- [25] B.R. Shreve, P.A. Moore, T.C. Daniel, D.R. Edwards, and D.M. Miller, Reduction of phosphorus in runoff from field-applied poultry litter using chemical amendments. *J. Environ. Qual.* Vol. 24, 1995, pp 106-111.