

Quantum Theory of Fields from Planck to Cosmic Scales

SIYAVASH H. SOHRAB

Robert McCormick School of Engineering and Applied Science

Department of Mechanical Engineering

Northwestern University, Evanston, Illinois 60208

s-sohrab@northwestern.edu

<http://www.mech.northwestern.edu/web/people/faculty/sohrab.php>

Abstract: - A scale invariant model of statistical mechanics is applied to describe a modified statistical theory of turbulence and its quantum mechanical foundations. Hierarchies of statistical fields from cosmic to *Planck* scales are described. Energy spectrum of equilibrium isotropic turbulence is shown to follow *Planck* law. Predicted velocity profiles of turbulent boundary layer over a flat plate at four consecutive scales of LED, LCD, LMD, and LAD are shown to be in close agreement with the experimental observations in the literature. The physical and quantum nature of time is described and a scale-invariant definition of time is presented and its relativistic behavior is examined. New paradigms for physical foundations of quantum mechanics as well as derivation of *Dirac* relativistic wave equation are introduced.

Key-Words: - Theory of turbulence; Quantum mechanics; Relativity; Dirac and Schrödinger equations; TOE.

1 Introduction

It is well known that the laws of nature appear to reveal ever increasing similarities over a broad range of scales of space and time from the exceedingly large scale of cosmology to the minute scale of quantum optics (Fig.1). The similarities between stochastic quantum fields [1-17] and classical hydrodynamic fields [18-29] resulted in recent introduction of a scale-invariant model of statistical mechanics [30], and its application to thermodynamics [31] and fluid mechanics [32].

More recently, the implication of the model to the statistical theory of turbulence [33, 34] was investigated. In the present study the physical foundations of the problems of turbulence and quantum mechanics are further examined. Homogenous isotropic turbulence is identified as a spectrum of eddies (energy levels) with *Gaussian* velocity distribution, *Planck* energy distribution, and *Maxwell-Boltzmann* speed distribution. The nature of dissipation spectrum of isotropic turbulence is examined and a derivation of *Dirac* relativistic wave equation is presented.

2 A Scale Invariant Model of Statistical Mechanics

Following the classical methods [35-39] the invariant definitions of density ρ_β , and velocity of *atom* \mathbf{u}_β , *element* \mathbf{v}_β , and *system* \mathbf{w}_β at the scale β are given as [31]

$$\rho_\beta = n_\beta m_\beta = m_\beta \int f_\beta d\mathbf{u}_\beta, \quad \mathbf{u}_\beta = \mathbf{v}_{\beta-1} \quad (1)$$

$$\mathbf{v}_\beta = \rho_\beta^{-1} m_\beta \int \mathbf{u}_\beta f_\beta d\mathbf{u}_\beta, \quad \mathbf{w}_\beta = \mathbf{v}_{\beta+1} \quad (2)$$

Similarly, the invariant definition of the peculiar and diffusion velocities are introduced as

$$\mathbf{V}'_\beta = \mathbf{u}_\beta - \mathbf{v}_\beta, \quad \mathbf{V}_\beta = \mathbf{v}_\beta - \mathbf{w}_\beta \quad (3)$$

such that

$$\mathbf{V}_\beta = \mathbf{V}'_{\beta+1} \quad (4)$$

For each statistical field, one defines particles that form the background fluid and are viewed as point-mass or "*atom*" of the field. Next, the *elements* of the field are defined as finite-sized composite entities each composed of an ensemble of "*atoms*" as shown in Fig.1.




Fig.1 A scale invariant view of statistical mechanics from cosmic to tachyon scales.

Finally, the ensemble of a large number of "elements" is defined as the statistical "system" at that particular scale.

3 Scale Invariant Forms of the Conservation Equations for Chemically Reactive Fields

Following the classical methods [35-37], the scale-invariant forms of mass, thermal energy, linear and angular momentum conservation equations at scale β are given as [40]

$$\frac{\partial \rho_{i\beta}}{\partial t_{\beta}} + \nabla \cdot (\rho_{i\beta} \mathbf{v}_{i\beta}) = \Omega_{i\beta} \quad (5)$$

$$\frac{\partial \varepsilon_{i\beta}}{\partial t_{\beta}} + \nabla \cdot (\varepsilon_{i\beta} \mathbf{v}_{i\beta}) = 0 \quad (6)$$

$$\frac{\partial \mathbf{p}_{i\beta}}{\partial t_{\beta}} + \nabla \cdot (\mathbf{p}_{i\beta} \mathbf{v}_{i\beta}) = -\nabla \cdot \mathbf{P}_{ij\beta} \quad (7)$$

$$\frac{\partial \boldsymbol{\pi}_{i\beta}}{\partial t_{\beta}} + \nabla \cdot (\boldsymbol{\pi}_{i\beta} \mathbf{v}_{i\beta}) = 0 \quad (8)$$

that involve the *volumetric density* of thermal energy $\varepsilon_{i\beta} = \rho_{i\beta} \tilde{h}_{i\beta}$, linear momentum $\mathbf{p}_{i\beta} = \rho_{i\beta} \mathbf{v}_{i\beta}$, and angular momentum $\boldsymbol{\pi}_{i\beta} = \rho_{i\beta} \boldsymbol{\omega}_{i\beta}$, $\boldsymbol{\omega}_{i\beta} = \nabla \times \mathbf{v}_{i\beta}$. Also, $\Omega_{i\beta}$ is the chemical reaction rate, $\tilde{h}_{i\beta}$ is the absolute enthalpy [40],

$$\tilde{h}_{i\beta} = \int_0^T c_{pi\beta} dT_{\beta} \quad (9a)$$

and $\mathbf{P}_{ij\beta}$ is the stress tensor [35]

$$\mathbf{P}_{ij\beta} = m_{\beta} \int (\mathbf{u}_{i\beta} - \mathbf{v}_{i\beta})(\mathbf{u}_{j\beta} - \mathbf{v}_{j\beta}) f_{\beta} d\mathbf{u}_{\beta} \quad (10)$$

In the derivation of (7) we have used the definition of the peculiar velocity (3) along with the identity

$$\overline{\mathbf{V}'_{i\beta} \mathbf{V}'_{j\beta}} = \overline{(\mathbf{u}_{i\beta} - \mathbf{v}_{i\beta})(\mathbf{u}_{j\beta} - \mathbf{v}_{j\beta})} = \overline{\mathbf{u}_{i\beta} \mathbf{u}_{j\beta}} - \mathbf{v}_{i\beta} \mathbf{v}_{j\beta} \quad (11)$$

The definition of absolute enthalpy in (9a) results in the definition of standard heat of formation $\tilde{h}_{i\beta}^{\circ}$ for chemical species i [40]

$$\tilde{h}_{i\beta}^{\circ} = \int_0^{T_0} c_{pi\beta} dT_{\beta} \quad (9b)$$

where T_0 is the standard temperature. The definition (9b) helps to avoid the conventional practice of arbitrarily setting the standard heat of formation of some species equal to zero. Furthermore, following

Nernst-Planck statement of the third law of thermodynamics one has $\tilde{h}_{\beta} \rightarrow 0$ in the limit $T_{\beta} \rightarrow 0$ as expected.

The classical definition of vorticity involves the curl of linear velocity $\nabla \times \mathbf{v}_{\beta} = \boldsymbol{\omega}_{\beta}$ thus giving rotational velocity of particle a secondary status in that it depends on its translational velocity \mathbf{v}_{β} . However, it is known that particle's rotation about its center of mass is independent of the translational motion of its center of mass. In other words, translational, rotational, and vibrational (pulsational) motions of particle are independent degrees of freedom that should not be necessarily coupled. To resolve this paradox, the iso-spin of particle at scale β is defined as the curl of the velocity at the next lower scale of $\beta-1$ [41]

$$\boldsymbol{\omega}_{\beta} = \nabla \times \mathbf{v}_{\beta-1} = \boldsymbol{\omega}_{\beta-1} = \nabla \times \mathbf{u}_{\beta} \quad (12)$$

such that the rotational velocity, while having a connection to some type of translational motion at internal scale $\beta-1$, retains its independent degree of freedom at the external scale β as desired. A schematic description of iso-spin and vorticity fields is shown in Fig.2. The nature of galactic vortices in cosmology and the associated dissipation have been discussed [25, 42].




Fig.2 Description of internal (iso-spin) versus external vorticity fields in cosmology [41].

The local velocity \mathbf{v}_{β} in (5)-(8) is expressed in terms of the convective \mathbf{w}_{β} and the diffusive \mathbf{V}_{β} velocities [40]

$$\mathbf{w}_{\beta} = \mathbf{v}_{\beta} - \mathbf{V}_{\beta g} \quad , \quad \mathbf{V}_{\beta g} = -D_{\beta} \nabla \ln(\rho_{\beta}) \quad (13a)$$

$$\mathbf{w}_{\beta} = \mathbf{v}_{\beta} - \mathbf{V}_{\beta tg} \quad , \quad \mathbf{V}_{\beta tg} = -\alpha_{\beta} \nabla \ln(\varepsilon_{\beta}) \quad (13b)$$

$$\mathbf{w}_{\beta} = \mathbf{v}_{\beta} - \mathbf{V}_{\beta hg} \quad , \quad \mathbf{V}_{\beta hg} = -v_{\beta} \nabla \ln(\mathbf{p}_{\beta}) \quad (13c)$$

$$\mathbf{w}_{\beta} = \mathbf{v}_{\beta} - \mathbf{V}_{\beta rhg} \quad , \quad \mathbf{V}_{\beta rhg} = -v_{\beta} \nabla \ln(\boldsymbol{\pi}_{\beta}) \quad (13d)$$

where $(\mathbf{V}_{\beta g}, \mathbf{V}_{\beta tg}, \mathbf{V}_{\beta hg}, \mathbf{V}_{\beta rhg})$ are respectively the diffusive, the thermo-diffusive, the translational and rotational hydro-diffusive velocities.

Because by definition fluids can only support compressive normal forces, following *Cauchy* the total stress tensor for fluids is expressed as [40]

$$\begin{aligned} \mathbf{P}_{ij\beta} &= p_\beta \delta_{ij\beta} - \lambda_\beta \nabla \cdot \mathbf{v}_\beta \delta_{ij} - 2\mu_\beta e_{ik} \delta_{ik} \delta_{ij} \\ &= p_\beta \delta_{ij} - \lambda_\beta \nabla \cdot \mathbf{v}_\beta \delta_{ij} - \mu_\beta \nabla \cdot \mathbf{v}_\beta \delta_{ij} \end{aligned} \quad (14)$$

Making the conventional *Stokes* assumption, i.e. setting the bulk viscosity b_β to zero, the two *Lame* constants will be related by [43]

$$b_\beta = \lambda_\beta + \frac{2}{3} \mu_\beta = 0 \quad (15)$$

and (14) reduces to [40]

$$\mathbf{P}_{ij\beta} = p_\beta \delta_{ij\beta} - \frac{1}{3} \mu_\beta \nabla \cdot \mathbf{v}_\beta \delta_{ij\beta} = (p_{i\beta} + p_{h\beta}) \delta_{ij\beta} \quad (16)$$

that involves thermodynamic $p_{i\beta}$ and hydrodynamic $p_{h\beta}$ pressures [40]. Following the classical methods [35-37], by substituting from (13)-(16) into (5)-(8) and neglecting cross-diffusion terms the invariant forms of conservation equations are written as [40]

$$\frac{\partial \rho_{i\beta}}{\partial t_\beta} + \mathbf{w}_\beta \cdot \nabla \rho_{i\beta} - D_{i\beta} \nabla^2 \rho_{i\beta} = \Omega_{i\beta} \quad (17)$$

$$\frac{\partial T_{i\beta}}{\partial t_\beta} + \mathbf{w}_\beta \cdot \nabla T_{i\beta} - \alpha_{i\beta} \nabla^2 T_{i\beta} = -\tilde{h}_{i\beta} \Omega_{i\beta} / (\rho_{i\beta} c_{pi\beta}) \quad (18)$$

$$\begin{aligned} \frac{\partial \mathbf{v}_{i\beta}}{\partial t_\beta} + \mathbf{w}_\beta \cdot \nabla \mathbf{v}_{i\beta} - \nu_{i\beta} \nabla^2 \mathbf{v}_{i\beta} &= -\frac{\nabla p_{i\beta}}{\rho_{i\beta}} \\ &+ \frac{1}{3} \nu_{i\beta} \nabla (\nabla \cdot \mathbf{v}_{i\beta}) - \frac{\mathbf{v}_{i\beta} \Omega_{i\beta}}{\rho_{i\beta}} \end{aligned} \quad (19)$$

$$\frac{\partial \boldsymbol{\omega}_{i\beta}}{\partial t_\beta} + \mathbf{w}_\beta \cdot \nabla \boldsymbol{\omega}_{i\beta} - \nu_{i\beta} \nabla^2 \boldsymbol{\omega}_{i\beta} = -\boldsymbol{\omega}_{i\beta} \cdot \nabla \mathbf{w}_\beta - \frac{\boldsymbol{\omega}_{i\beta} \Omega_{i\beta}}{\rho_{i\beta}} \quad (20)$$

The main new feature of the modified form of the equation of motion (19) is its linearity due to the difference between the convective \mathbf{w}_β versus the local velocity \mathbf{v}_β as compared with the non-linear classical *Navier-Stokes* equation of motion

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} - \nu \nabla^2 \mathbf{v} = -\frac{\nabla p}{\rho} + \frac{1}{3} \nu \nabla (\nabla \cdot \mathbf{v}) \quad (21)$$

The convection velocity in (19) must be obtained from the solution of the equation of motion for outer potential flow at the next larger scale $\mathbf{w}_\beta = \mathbf{v}_{\beta+1}$. One further notes that in (19) the absence of convection results in none-homogeneous diffusion equation while in (21) the absence of velocity leads to the vanishing of almost the entire equation of motion [40]. Also, the linearity of (19), in harmony with *Carrier* equation [44], resolves the classical paradox of drag reciprocity [45, 46].

The central question concerning *Cauchy* equation of motion (7) say at $\beta = m$ is how many “molecules” are included in the definition of the mean molecular velocity $\mathbf{v}_{mj} = \langle \mathbf{u}_{mj} \rangle = \mathbf{u}_{cj}$. One can identify three distinguishable cases: (1) When $\mathbf{v}_m = \mathbf{u}_c$ is itself random then all three velocities ($\mathbf{u}_m, \mathbf{v}_m, \mathbf{V}'_m$) in (3) are random in a stochastically stationary field made of ensembles of clusters and molecules with *Brownian* motions and hence *Gaussian* velocity distribution, *Planck* energy distribution, and *Maxwell-Boltzmann* speed distribution. If in addition both the vorticity $\boldsymbol{\omega}_m = \nabla \times \mathbf{v}_m = 0$ as well as the iso-spin (12) $\boldsymbol{\omega}_m = \nabla \times \mathbf{u}_m = 0$ are zero, then for an incompressible flow the continuity equation (5) and *Cauchy* equation of motion (7) lead to *Bernoulli* equation. In Section 7 it will be shown that under the above mentioned conditions *Schrödinger* equation can be directly derived from *Bernoulli* equation such that the energy spectrum of the equilibrium field will be governed by quantum mechanics and hence by *Planck* law. (2) When $\mathbf{v}_m = \mathbf{u}_c$ is not random but the vorticity vanishes $\boldsymbol{\omega}_m = 0$ the flow is irrotational and ideal, inviscid $\mu_m = 0$, and once again one obtains *Bernoulli* equation from (5) and (7) with the solution given by the classical potential flow. (3) When $\mathbf{v}_m = \mathbf{u}_c$ is not random and vorticity does not vanish $\nabla \times \mathbf{v}_m \neq 0$, the rotational non-ideal viscous $\mu_m \neq 0$ flow will be governed by the equation of motion (19) with the convection velocity $\mathbf{w}_\beta = \mathbf{v}_{\beta+1}$ obtained from the solution of potential flow at the next larger scale of $\beta+1$. In Section 9 it will be shown that the viscous equation of motion (19) is associated with *Dirac* relativistic wave equation. In the sequel, all three cases of flow conditions discussed above will be examined.

4 A Modified Statistical Theory of Turbulence

The invariant model of statistical mechanics (1)-(4) suggests that all statistical fields shown in Fig.1 are turbulent fields and governed by (5)-(8) [33, 34]. First, let us start with the field of laminar molecular dynamics LMD when molecules, clusters of molecules (cluster), and cluster of clusters of molecules (eddy) form the “atom”, the “element”, and the “system” with the velocities ($\mathbf{u}_m, \mathbf{v}_m, \mathbf{w}_m$). Similarly, the fields of laminar cluster-dynamics LCD and eddy-dynamics

LED will have the velocities ($\mathbf{u}_e, \mathbf{v}_e, \mathbf{w}_e$), and ($\mathbf{u}_e, \mathbf{v}_e, \mathbf{w}_e$) in accordance with (1)-(2). For the fields of LED, LCD, and LMD, typical characteristic “atom”, element, and system lengths are

$$\text{EED } (\ell_e, \lambda_e, L_e) = (10^{-5}, 10^{-3}, 10^{-1}) \text{ m} \quad (22a)$$

$$\text{ECD } (\ell_e, \lambda_e, L_e) = (10^{-7}, 10^{-5}, 10^{-3}) \text{ m} \quad (22b)$$

$$\text{EMD } (\ell_m, \lambda_m, L_m) = (10^{-9}, 10^{-7}, 10^{-5}) \text{ m} \quad (22c)$$

If one applies the same (atom, element, system) = ($\ell_\beta, \lambda_\beta, L_\beta$) relative sizes in (22) to the entire spatial scale of Fig.1, then the resulting cascades or hierarchy of overlapping statistical fields will appear as schematically shown in Fig.3.




Fig.3 Hierarchy of statistical fields with ($\ell_\beta, \lambda_\beta, L_\beta$) from cosmic to Planck scales [34].

According to Fig.3, starting from the hydrodynamic scale ($10^3, 10^1, 10^{-1}, 10^{-3}$) after seven generations of statistical fields one reaches the electro-dynamic scale with the element size 10^{-17} , and exactly after seven more generations one reaches *Planck* length scale $(\hbar G / c^3)^{1/2} \approx 10^{-35} \text{ m}$, where G is the gravitational constant. Similarly, seven generations of statistical fields separate the hydrodynamic scale ($10^3, 10^1, 10^{-1}, 10^{-3}$) from the scale of planetary dynamics (astrophysics) 10^{17} and the latter from galactic-dynamics (cosmology) 10^{35} m .

The left hand side of Fig.1 corresponds to equilibrium statistical fields when the velocities of elements of the field are random since at thermodynamic equilibrium particles i.e. oscillators of such statistical fields will have normal or *Gaussian* velocity distribution. For example, for

stationary homogeneous isotropic turbulence at EED scale, the experimental data of *Townsend* [47] confirms the *Gaussian* velocity distribution of eddies as shown in Fig.4.




Fig.4 Measured velocity distribution in isotropic turbulent flow [47].

Because at thermodynamic equilibrium the mean velocity of each particle, *Heisenberg-Kramers* virtual oscillator [48], vanishes $\langle \mathbf{u}_\beta \rangle = 0$ the energy of particle oscillating in two directions ($x+$, $x-$) is expressed as

$$\begin{aligned} \varepsilon_\beta &= m_\beta \langle u_{\beta x+}^2 \rangle / 2 + m_\beta \langle u_{\beta x-}^2 \rangle / 2 \\ &= m_\beta \langle u_{\beta x+}^2 \rangle = \langle p_\beta \rangle \langle \lambda_\beta^2 \rangle^{1/2} \langle v_\beta^2 \rangle^{1/2} \end{aligned} \quad (23)$$

where $m_\beta \langle u_{\beta x+}^2 \rangle^{1/2} = \langle p_\beta \rangle$ is the root-mean-square momentum of particle and $\langle u_{\beta x+}^2 \rangle = \langle u_{\beta x-}^2 \rangle$ by equipartition principle. At any scale β , the result (23) can be expressed in terms of either frequency or wavelength

$$\varepsilon_\beta = m_\beta \langle u_\beta^2 \rangle = \langle p_\beta \rangle \langle \lambda_\beta^2 \rangle^{1/2} \langle v_\beta^2 \rangle^{1/2} = h_\beta v_\beta \quad (24)$$

$$\varepsilon_\beta = m_\beta \langle u_\beta^2 \rangle = \langle p_\beta \rangle \langle v_\beta^2 \rangle^{1/2} \langle \lambda_\beta^2 \rangle^{1/2} = k_\beta \lambda_\beta \quad (25)$$

when the definition of stochastic *Planck* and *Boltzmann* factors are introduced as [33]

$$h_\beta = \langle p_\beta \rangle \langle \lambda_\beta^2 \rangle^{1/2} \quad (26)$$

$$k_\beta = \langle p_\beta \rangle \langle v_\beta^2 \rangle^{1/2} \quad (27)$$

At the important scale of EKD (Fig.1) corresponding to *Casimir* vacuum [49] composed of photon gas, the universal constants of *Planck* [50, 51] and *Boltzmann* [31] are obtained from (24)-(25) as

$$h = h_k = m_k c \langle \lambda_k^2 \rangle^{1/2} = 6.626 \times 10^{-34} \text{ J-s} \quad (28)$$

$$k = k_k = m_k c \langle v_k^2 \rangle^{1/2} = 1.381 \times 10^{-23} \text{ J/K} \quad (29)$$

Next, following *de Broglie* hypothesis for the wavelength of matter waves [2]

$$\lambda_\beta = h / p_\beta \quad (30)$$

the frequency of matter waves is defined as [31]

$$v_{\beta} = k / p_{\beta} \quad (31)$$

When matter and radiation are in the state of thermodynamic equilibrium (30) and (31) can be expressed as

$$h_{\beta} = h_k = h, \quad k_{\beta} = k_k = k \quad (32)$$

The definitions (28) and (29) result in the gravitational mass of photon [31]

$$m_k = (hk / c^3)^{1/2} = 1.84278 \times 10^{-41} \text{ kg} \quad (33)$$

that is much larger than the reported value of $4 \times 10^{-51} \text{ kg}$ [52]. The finite gravitational mass of photons was anticipated by *Newton* [53] and is in accordance with the *Einstein-de Broglie* theory of light [54-58]. *Avogadro-Loschmidt* number was predicted as [31]

$$N^0 = 1/(m_k c^2) = 6.0376 \times 10^{23} \quad (34)$$

leading to the modified value of the universal gas constant

$$R^0 = N^0 k = 8.338 \text{ kJ/(kmol-K)} \quad (35)$$

The classical definition of thermodynamic temperature that is based on two degrees of freedom

$$kT' = m \langle u_x^2 \rangle = 2m \langle u_{x+}^2 \rangle \quad (36)$$

was recently modified to a new definition based on single degree of freedom [59]

$$kT = m \langle u_{x+}^2 \rangle \quad (37)$$

such that

$$T' = 2T \quad (38)$$

The factor 2 in (38) results in the predicted speed of sound in nitrogen [60]

$$a = \langle u_{x+}^2 \rangle^{1/2} = \sqrt{p / (2\rho)} \approx 357 \text{ m/s} \quad (39)$$

($a \approx 350 \text{ m/s}$ for air) in close agreement with observations. Therefore, the square root of 2 in (39) resolves the classical problem of *Newton* concerning his prediction of velocity of sound as

$$a = \sqrt{p / \rho} \quad (40)$$

discussed by *Chandrasekhar* [61]

“Newton must have been baffled, not to say disappointed. Search as he might, he could find no flaw in his theoretical framework—neither could Euler, Lagrange, and Laplace; nor, indeed, anyone down to the present”

The factor of 2 in (38) also leads to the modified value of the mechanical equivalent of heat J [59]

$$J = 2J_c = 2 \times 4.169 = 8338 \text{ Joules/(kcal)} \quad (41)$$

where the value $J_c = 4.169 \approx 4.17 \text{ [kJ/kcal]}$ is the average of the two values $J_c = (4.15, 4.19)$ reported by *Pauli* [62]. The number in (41) is thus identified as the universal gas constant (35) when expressed in appropriate MKS system of units

$$R^0 = kN^0 = 8338 \text{ Joules/(kmol-K)} \quad (42)$$

The modified value of the universal gas constant (42) was recently identified [63] as *De Pretto* number 8338 that appeared in the mass–energy equivalence equation of *De Pretto* [64]

$$E = mc^2 \text{ Joules} = mc^2 / 8338 \text{ kcal} \quad (43)$$

5 Energy Spectra of Isotropic Turbulence given by Planck Energy Distribution Law

The field of isotropic homogeneous turbulence is identified as equilibrium eddy dynamics EED, Fig.1, with turbulent eddies defined as clusters of molecular clusters constituting the elements of the field. In a recent investigation [33], it was shown that the energy spectrum of eddies in isotropic turbulence is governed by invariant *Planck* energy distribution law [33, 50]

$$\frac{\epsilon_{\beta} dN_{\beta}}{V} = \frac{8\pi h}{u_{\beta}^3} \frac{v_{\beta}^3}{e^{h v_{\beta} / kT} - 1} dv_{\beta} \quad (44)$$

schematically shown in Fig.5.




Fig.5 Planck energy distribution law governing the energy spectrum of eddies at the temperature $T = 300 \text{ K}$.

From application of *Boltzmann* distribution of molecules in clusters and clusters in eddies it was found that [33]

$$N_{\beta} = \frac{g_{\beta}}{e^{h\nu_{\beta}/kT} - 1} \quad (45)$$

that along with *Rayleigh-Jeans* number for degeneracy [51, 65, 66]

$$dg_{\beta} = \frac{8\pi V}{u_{\beta}^3} v_{\beta}^2 dv_{\beta} \quad (46)$$

and $\varepsilon_{\beta} = h\nu_{\beta}$ give (44). It is interesting to examine a new perspective of (46) that is related to determination of number of degeneracy from field quantization [67]. The number of degeneracy of particles, *Heisenberg-Kramers* virtual oscillators [48], in spherical volume V_S is written as

$$g_{\beta} = 2V_S / \lambda_{\beta}^3 \quad (47)$$

where λ_{β}^3 is the *rectangular* volume occupied by each oscillator $V_o = \lambda_{\beta}^3$ when due to isotropy $\lambda_{\beta} = \langle \lambda_{x\beta}^2 \rangle^{1/2} = \langle \lambda_{y\beta}^2 \rangle^{1/2} = \langle \lambda_{z\beta}^2 \rangle^{1/2}$ and the factor 2 comes from allowing particles to have two modes either (up) or (down) iso-spin (polarization). The system spherical V_S and rectangular V volumes are related as

$$V_S = \frac{4\pi}{3} R^3 = \frac{4\pi}{3} L^3 = \frac{4\pi}{3} V \quad (48)$$

For systems in thermodynamic equilibrium the temperature $3kT_{\beta} = m_{\beta} \langle u_{\beta}^2 \rangle$ will be constant and hence $\langle u_{\beta}^2 \rangle^{1/2} = \langle \lambda_{\beta}^2 \rangle^{1/2} \langle v_{\beta}^2 \rangle^{1/2}$ or

$$\lambda_{\beta} = u_{\beta} / v_{\beta} \quad (49)$$

Substituting from (48) and (49) into (47) results in

$$g_{\beta} = \frac{8\pi V}{3u_{\beta}^3} v_{\beta}^3 \quad (50)$$

that leads to the number of oscillators between frequencies ν_{β} and $\nu_{\beta} + d\nu_{\beta}$

$$dg_{\beta} = \frac{8\pi V}{u_{\beta}^3} v_{\beta}^2 dv_{\beta} \quad (51)$$

in accordance with (46).

With *Gaussian* velocity distribution, Fig.4, the same chain of reasoning as employed in the classical kinetic theory of gas requires that the distribution of the speeds of oscillators (eddies) in stationary isotropic turbulence be given by the invariant *Maxwell-Boltzmann* distribution function

$$\frac{dN_{u_{\beta}}}{N} = 4\pi \left(\frac{m_{\beta}}{2\pi kT_{\beta}} \right)^{3/2} u_{\beta}^2 e^{-m_{\beta} u_{\beta}^2 / 2kT_{\beta}} du_{\beta} \quad (52)$$

By (52), one arrives at a hierarchy of embedded *Maxwell-Boltzmann* distribution functions for EED, ECD, and EMD scales shown in Fig.6.




Fig.6 Maxwell-Boltzmann speed distribution viewed as stationary spectra of cluster sizes for EED, ECD, and EMD scales at 300 K [33].

To summarize, at equilibrium the statistical fields of any scale shown on the left-hand-side of Fig.1 will have elements with (a) *Gaussian* velocity distribution, (b) *Planck* energy distribution, and (c) *Maxwell-Boltzmann* speed distribution. For the conventional fields of ECD and EMD, it is well known that the conditions (a) and (c) are true and because of equilibrium between matter and radiation fields it is reasonable to expect that the condition (b) above will also apply to LCD and LMD. At the scale of LED, the conditions (a) shown by *Townsend's* data in Fig.4 and hence condition (c) are known to hold. Preliminary examination of the three-dimensional energy spectrum $E(k)$ for isotropic turbulence measured by *Van Atta and Chen* [68] in Fig.7




Fig.7 Normalized three-dimensional energy spectra for isotropic turbulence [68].

appears to support the validity of condition (b).

The schematic diagram given in Fig.8 from *Landahl and Mollo-Christensen* [69]




Fig.8 Behavior of three-dimensional spectrum $E(k, t)$ in various wave number ranges [69].

also appears to support the validity of condition (b) namely that *Planck* law governs the energy spectrum of eddies in isotropic turbulence. Unfortunately, in spite of the large number of experimental studies, data that clearly and directly show the variation of three-dimensional energy spectrum $E(k)$ with wave number k are still lacking.

According to Fig.8, the *Kolmogorov-Obukhov* $k^{-5/3}$ law [22, 23] is a local feature, valid only in the inertial subrange, of the more universal *Planck* law (44). For stationary isotropic turbulent fields, energy input into the system cascades down to smaller and smaller oscillators until it finally reaches the *Kolmogorov* dissipation length scale η_k at which point all of the added energy blends into background white noise and is removed from the system as heat. In view of Fig.6, η_k is naturally identified as the atomic length ℓ_e of LED scale which is the same as the element λ_c of LCD scale

$$\eta_k = \ell_e = \lambda_c \quad (53)$$

that appears in *Boussinesq* eddy diffusivity [70]

$$\nu_e = \frac{1}{3} \ell_e u_e = \frac{1}{3} \lambda_c v_c \quad (54)$$

On the other hand, the kinematic viscosity of LCD field is related to the “atomic” length ℓ_c or the molecular mean free path λ_m that appears in *Maxwell’s* formula for kinematic viscosity [30]

$$\nu_c = \frac{1}{3} \ell_c u_c = \frac{1}{3} \lambda_m v_m, \quad (55)$$

associated with viscous dissipation in fluid mechanics.

In stationary isotropic turbulent fields, energy flux occurs between randomly moving eddies of diverse size while leaving the system stochastically stationary in time. A schematic diagram of energy

flux across hierarchies of eddies from large to small size is shown in Fig.9 from the study by *Lumley et al* [71]




Fig. 9 A realistic view of spectral energy flux [71].

In the following section, it will be suggested that the exchange of particles, clusters, between various size eddies (energy levels) is governed by quantum mechanics through an invariant *Schrödinger* equation. Hence, the stochastically stationary state of various size eddies will be parallel to *Bohr’s stationary states* in atomic theory [48].

For stationary isotropic turbulent fields the net energy flux across hierarchies of eddies should be a constant independent of viscosity such as is commonly assumed in the inertial subrange. Also, at equilibrium, the temperature will be constant $3k_B T = m \langle u^2 \rangle = \varepsilon$. Therefore, if one expresses the cluster energy as $\varepsilon = mu^2$, and the eddy energy as $E = \sum \varepsilon = \alpha \varepsilon$, with the cluster velocity given as $u = v\lambda = 2\pi v/\kappa$ such that $du = d(2\pi v/\kappa) \propto \kappa^{-2} d\kappa$ where $\kappa = 2\pi/\lambda$ is the wave number, one arrives at

$$\begin{aligned} dE &= F d\kappa = 2\alpha m u du \propto \alpha \varepsilon^{2/3} (u \varepsilon^{-2/3}) \kappa^{-2} d\kappa \\ &= \alpha \varepsilon^{2/3} (\kappa^{-1} \kappa^{4/3}) \kappa^{-2} d\kappa = \alpha \varepsilon^{2/3} \kappa^{-5/3} d\kappa \end{aligned} \quad (56)$$

leading to the distribution function

$$F(\kappa) = \alpha \varepsilon^{2/3} \kappa^{-5/3} \quad (57)$$

that is the *Kolmogorov-Obukhov* $\kappa^{-5/3}$ law [22, 23].

The most central concept associated with turbulent dissipation is the spectral definition of turbulent viscosity introduced by *Heisenberg* [26]

$$\nu_k = \kappa \int_{\kappa}^{\infty} \left(\sqrt{\frac{F(\kappa'')}{\kappa'^3}} \right) d\kappa'' \quad (58)$$

This is because the spectral definition of kinematic viscosity (58) suggests that in stationary isotropic turbulence the dissipation spectrum should be closely related to the energy spectrum and thereby to *Planck* energy spectrum. Preliminary examination of dissipation spectrum shown in Fig. 10 from *McComb and Shanmugasundaram* [72] appears to support such a conjecture.




Fig.10 Comparison of scaled one-dimensional dissipation spectrum with experimental data [72].

The experimental data of Fig.10 along with *Planck* distribution function as well as this same distribution function shifted by a constant amount of energy are shown in Fig.11. One may view the experimental data in Fig.10 as dissipation spectrum of energy associated with the *Planck* energy spectrum plus a constant amount of energy added in order to maintain the turbulent field stationary.




Fig.11 One-dimensional dissipation spectrum [72] compared with (1) Planck energy distribution (2) Planck energy distribution with constant displacement.

Similar comparison with *Planck* distribution as shown in Fig.11 is obtained with the more recent experimental data of one-dimensional dissipation spectrum of isotropic turbulence from study by *Saddoughi and Veeravalli* [73] given in Fig.12.




Fig.12 One-dimensional dissipation spectra measured (a) u_1 -spectrum (b) u_2 -spectrum (c) u_3 -spectrum [73].

In another very recent experimental investigation the energy spectra of turbulent flow within the boundary layer in close vicinity of rigid wall was measured by *Marusic et al.*, [74] and the reported energy spectra shown in Fig.(13) appear to have profiles quite similar to *Planck* distribution law.




Fig.13 Reynolds number evolution of the premultiplied energy spectra of streamwise velocity at the inner-peak location ($z^+ = 15$) for the true measurements (A) and the prediction based on the filtered u signal measured in the log region (B) [74].

The normalized three-dimensional energy spectrum for homogeneous isotropic turbulent field was obtained from the transformation of one-dimensional spectrum of *Lin* [75] by *Ling and Huang* [76] as

$$E^* = \frac{\alpha^2}{3} (K^* + \alpha K^{*2}) \exp(-\alpha K^*) \quad (59)$$

with the distribution shown in Fig.14.




Fig.14 Normalized three-dimensional energy spectrum of weak turbulence [76].

The similarity between Fig.14 and the *Planck* distribution function shown in Fig.5 is apparent and with $h/kT = c_1$ one can express (43) as

$$\varepsilon_v dN = c_2 v^3 \exp(-c_1 v) / [1 - \exp(-c_1 v)] \quad (60)$$

to facilitate comparison with (59). Further future investigations on systematic comparisons between measurements of three-dimensional energy spectrum of equilibrium isotropic turbulence and *Planck* distribution (Fig.5) are needed.

Because the velocity distribution of eddies in isotropic turbulence is known to be *Gaussian*, Fig.4, the distribution of the speed of eddies in isotropic

turbulence must follow *Maxwell-Boltzmann* distribution function in accordance with the kinetic theory of gas. On the other hand, in a recent investigation [33] the invariant *Maxwell-Boltzmann* distribution function was directly derived from the invariant *Planck* energy distribution function. Therefore, it is expected that the energy spectrum of eddies in isotropic turbulence should follow the *Planck* law [33, 34].

In the scale invariant statistical theory of turbulence described above the statistical nature of the problem does not arise from the notion of fluid instabilities as in the classical theories of turbulence, but rather arises from similar considerations as those in the kinetic theory of gas in harmony with perceptions of *Heisenberg* [26]

“Turbulence is an essentially statistical problem of the same type as one meets in statistical mechanics, since it is the problem of distribution of energy among a very large number of degrees of freedom. Just as in Maxwell theory this problem can be solved without going into details of the mechanical motions, so it can be solved here by simple considerations of similarity.”

It was also emphasized by *Heisenberg* that the study of flow instabilities cannot in principle lead to the understanding of turbulent phenomena itself as noted by *Chandrasekhar* [77]

“However, as Heisenberg has recently emphasized, investigation of stability along these lines, even if successful, cannot, in principle, lead to an understanding of the phenomena of turbulence itself; for the basic problem of turbulence is of an entirely different character. That this is the case becomes apparent when we ask ourselves the very elementary question, “What is the reason that a phenomena like turbulence can occur at all?”. The answer must be that an ideal fluid is a mechanical system with very large number of degrees of freedom and that, in consequence, it is theoretically capable of a very large number of different types of motions. Laminar flow is only one of the many possible motions that the system is capable of, and to expect that it will always be realized is as futile as to expect that in a gas we shall find all the molecules moving with the same velocity parallel to one another. It is far more likely that all forms of possible motions will be simultaneously present. The fundamental problem of turbulence would therefore appear to be a statistical one of specifying the probability with which the various types of motions may occur and are present. Stated in this way, it is clear that the problem of turbulence has an analogy with the problem of analyzing a continuous spectrum of radiation.”

The significance of analogy between the energy spectra in turbulence and in optics, dry hydrodynamics [17], was further emphasized by *Chandrasekhar* [77]

“Now, returning to the optical analogy I referred to earlier, we know that under condition of equilibrium the distribution of energy in the continuous spectrum will be that given by Planck’s law. We may ask whether a similar equilibrium spectrum exists for turbulence. In answering this question, we must keep in mind one important distinction between the optical analogue and turbulence. In optical case the equilibrium Planck spectrum will be reached, no matter what the initial distribution is. In contrast, turbulence can be maintained only by external energy, like continuous stirring, the energy available from thermal instability, or rotation in differentially rotating atmosphere. In other words, energy is required for the maintenance of turbulence; in the absence of such an agency, turbulence will decay and the spectrum will be a function of time.”

The optical analogy discussed by *Chandrasekhar* in the above quotation becomes complete if one allows a constant energy input at small wave numbers (at system scale L_e of EED) that moves through the hierarchy of eddies (Fig.9) until it is dissipated into heat that is removed at *Kolmogorov* scale $\eta_k = \ell_e$.

According to the modified statistical theory of turbulence [33, 34], and in harmony with the perceptions of *Heisenberg* and *Chandrasekhar* discussed above, all flows that may appear laminar at scale β are actually bulk advection of turbulent flows at the scale of $\beta-1$. The hierarchies of embedded turbulent flows are most clearly seen in turbulent boundary layer over a flat plate when the solutions of (19) at outer $\beta+1$ and inner β scales were respectively found to be [34]

$$v_{\beta+1}^+ = 5 + 8(2/\sqrt{\pi})^2 \operatorname{erf}(y_{\beta-1}^+/32) \quad (61)$$

and

$$v_{\beta}^+ = 8(2/\sqrt{\pi}) \operatorname{erf}(y_{\beta-1}^+/8) \quad (62)$$

For example, the solutions in (61) and (62) at $\beta+1 = e$ and $\beta = c$ correspond to LED and LCD and their comparisons with experimental data [36, 69, 78-80] are shown in Fig.15.




Fig.15 Comparison between the predicted velocity profiles from (61) and (62) at LED and LCD scales and experimental data in the literature [36, 69, 78-80].

Similarly, the solutions in (61) and (62) at $\beta+1 = c$ and $\beta = m$ correspond to LCD and LMD and at $\beta+1 = m$ and $\beta = a$ corresponding to LMD and LAD and their comparisons with experimental data of *Lancien et al.* [81] and *Meinhart et al.* [82] are shown in Figs.16 and 17, respectively.




Fig.16 Comparisons between the predicted velocity profiles from (61) and (62) at LCD and LMD scales and experimental data in the literature [81].




Fig.17 Comparisons between the predicted velocity profiles from (61) and (62) at LMD and LAD scales and experimental data in the literature [82]

The results in Figs.15-17 show close agreement between predictions and measurements for four consecutive statistical fields of LED, LCD, LMD, and LAD spanning a factor of 10^8 in spatial range.

An important observation concerning the results shown in Fig.15-17 is the absence of any transition region between the velocity fields of adjacent scales. This is to be expected since both flow fields at scales β and $\beta-1$ are turbulent and all that is needed in connecting the two solutions is the proper “renormalization” of the coordinates as described in [34]. While the outer most turbulent field experiences the uniform free stream velocity w_β , all subsequent inner velocity fields match the slope $(dv^+/dy^+)_\beta$ of the corresponding outer flow and hence must match the linear velocity $v_\beta^+ = y_\beta^+$.

6 The Natures of Potential and Viscous Flows

In this section, the natures of potential flow and viscous boundary layer flow will be examined by application of *Cauchy* (7) and the modified form of equation of motion (19), respectively. This is in

part motivated by the fact that potential flow fields are needed for the derivation of *Schrödinger* equation to be discussed in the following section. For convenience, the well known problem of two-dimensional stagnation point flow will be considered. For potential flows the velocity may be expressed in terms of the velocity potential Φ_β as

$$\mathbf{v}_\beta = -\nabla \Phi_\beta \quad (63)$$

Except for a few [43, 83], the minus sign in (63) is neglected in most textbooks on fluid mechanics including some classical texts [27, 28]. However, it is emphasized here that the choice of minus sign in (63) is important because it insures that the fluid flows in the direction of decreasing potential in harmony with *Ohm's* law for current density

$$\mathbf{J} = \rho_e \mathbf{V}_e = -\sigma_e \nabla \Phi_e \quad (64)$$

that governs the diffusional flux (drift) of electrons through conductors in electrodynamics.

For the classical problem of two-dimensional stagnation point flow the complex potential

$$Z_\beta(z) = \Phi_\beta + i\Psi_\beta = z^2/2 = (x^2 - y^2)/2 + ixy \quad (65)$$

with $z = x + iy$ leads to the well known velocity potential and stream function

$$\Phi_\beta = (x^2 - y^2)/2 \quad (66)$$

$$\Psi_\beta = xy \quad (67)$$

with some of the streamlines shown in Fig.18.




Fig.18 Streamlines and iso-potential lines for stagnation flow.

The *Cauchy-Riemann* conditions and the velocity field from (63), (66), and (67) are

$$-\frac{\partial \Phi_\beta}{\partial x} = -\frac{\partial \Psi_\beta}{\partial y} = v_x = -x \quad (68)$$

$$-\frac{\partial \Phi_\beta}{\partial y} = \frac{\partial \Psi_\beta}{\partial x} = v_y = y \quad (69)$$

One notes that the velocity (68)-(69) that is perpendicular to the equi-potential lines is in the direction of decreasing potential as shown in Fig.18. For flow in the reverse direction the sign of the complex potential (65) should be changed such that $Z_\beta(z) = -z^2/2$.

Next, for the study of viscous flow within the thin boundary layer at the lower scale $\beta-1$ let us consider the reverse of the outer potential flow (68)-(69) that leads to dimensionless convective velocity of the inner viscous layer [84]

$$v_{x\beta} = w_{x\beta-1} = x, \quad v_{y\beta} = w_{y\beta-1} = -y \quad (70)$$

With the conventional assumption of negligible pressure gradient across boundary layer and $\partial v_x / \partial x' \ll \partial v_x / \partial y'$, $\partial^2 / \partial x'^2 \ll \partial^2 / \partial y'^2$ one obtains for an incompressible flow from (19)

$$w_y \frac{\partial v_x}{\partial y} = \frac{\partial^2 v_x}{\partial y^2} \quad (71)$$

that with the boundary conditions

$$v_x(0) = 0, \quad v_x(\infty) = w_x = x \quad (72)$$

and upon substitution from (70) leads to the solution

$$v_x = x \operatorname{erf}(y/\sqrt{2}) \quad (73)$$

Also, from the continuity equation (5) for an incompressible flow one obtains

$$v_y = -\int_0^y \operatorname{erf}(y/\sqrt{2}) dy \quad (74)$$

leading to the stream function

$$\Psi = x \int \operatorname{erf}(y/\sqrt{2}) dy \quad (75)$$

with some calculated streamlines shown in Fig.19.




Fig.19 Streamlines in the boundary layer of stagnation point flow from (75).

7 Invariant Schrödinger Equation and Quantum Mechanical Foundations of Turbulence

The fact that the energy spectrum of equilibrium isotropic turbulence is given by *Planck* distribution (Figs.5, 9, 13, 14) is a strong evidence for quantum mechanical foundation of turbulence [33, 34]. This is further supported by recent derivation of invariant *Schrödinger* equation from invariant *Bernoulli* equation [33] to be further examined in the present section. As stated in Sec.3, when the three velocities ($\mathbf{u}_m, \mathbf{v}_m, \mathbf{v}'_m$) in (3) are all random the system is composed of an ensemble of clusters and molecules under equilibrium state. When both vorticity and iso-spin (12) are zero and one considers non-reactive incompressible fluid, the flow field will be potential and the continuity (5) and *Cauchy* equation of motion (7) lead to invariant *Bernoulli* equation that upon substitution from (63) gives [33]

$$\frac{-\partial(\rho_\beta \Phi_\beta)}{\partial t} + \frac{[-\nabla(\rho_\beta \Phi_\beta)]^2}{2\rho_\beta} + p_\beta = \text{const} \quad t=0 \quad (76)$$

The constant in (76) is set to zero since pressure to be identified as potential is only defined to within an additive constant. Comparison of (76) with the *Hamilton-Jacobi* equation of classical mechanics [2]

$$\frac{\partial S}{\partial t} + \frac{(\nabla S)^2}{2m} + U = 0 \quad (77)$$

results in introduction of the invariant action [33, 85]

$$S_\beta(\mathbf{x}, t) = -\rho_\beta \Phi_\beta \quad (78)$$

By the results in (63) the gradient of the action (78) becomes momentum in accordance with the classical results [3]

$$\nabla S_\beta(\mathbf{x}, t) = -\rho_\beta \nabla \Phi_\beta = \rho_\beta \mathbf{v}_\beta = \mathbf{p}_\beta \quad (79)$$

In addition to absence of vorticity $\nabla \times \mathbf{v}_\beta = 0$ required for potential flow one further requires the iso-spin (12) to also vanish

$$\nabla \times \mathbf{u}_\beta = 0, \quad \mathbf{u}_\beta = -\nabla \Phi_{o\beta} \quad (80)$$

such that from (3) one obtains

$$\nabla \times \mathbf{V}'_\beta = 0, \quad \mathbf{V}'_\beta = -\nabla \Phi'_\beta \quad (81)$$

Next, one considers the peculiar velocity \mathbf{V}'_β in (3) as a small perturbation

$$\mathbf{u}_\beta - \mathbf{v}_\beta = \varepsilon \mathbf{V}'_\beta \quad (82)$$

and in view of (63), substitutions from (78)-(81) into (82) give

$$\rho_\beta \Phi_\beta = \rho_\beta \Phi_{o\beta} - \varepsilon \rho_\beta \Phi'_\beta, \quad \varepsilon \ll 1 \quad (83)$$

By (78), and (80)-(83) one arrives at

$$S_\beta = S_{o\beta} - \varepsilon S'_\beta = S_{o\beta} - \varepsilon \Psi_\beta \quad (84)$$

where $S_{o\beta} = -\rho_\beta \Phi_{o\beta}$, and the *wave function* Ψ_β of quantum mechanics is defined as [33, 85]

$$\Psi_\beta = S'_\beta = -\rho_\beta \Phi'_\beta \quad (85)$$

Substituting from (84) into (76) and separating terms of equal powers of ε leads to

$$\frac{\partial S_{o\beta}}{\partial t} + \frac{(\nabla S_{o\beta})^2}{2\rho_\beta} + \tilde{U}_\beta = 0 \quad (86)$$

$$\frac{\partial \Psi_\beta}{\partial t} + \frac{\nabla S_{o\beta} \nabla \Psi_\beta}{\rho_\beta} = 0 \quad (87)$$

where the volumetric potential energy density is defined as [28]

$$\tilde{U}_\beta \equiv p_\beta = n_\beta \bar{U}_\beta \quad (88)$$

To analyze (86), one introduces the moving coordinate

$$Z_\beta = \mathbf{x}_\beta - \mathbf{u}_\beta t_\beta \quad (89)$$

such that

$$\nabla_x S_{o\beta} = \nabla_z S_{o\beta}, \quad \partial S_{o\beta} / \partial t = -\mathbf{u}_\beta \nabla_z S_{o\beta} \quad (90)$$

Substitutions from (88), (89) and (90) in (86) give the conservation of energy

$$\tilde{E}_\beta = \tilde{T}_\beta + \tilde{U}_\beta, \quad \bar{E}_\beta = \bar{T}_\beta + \bar{U}_\beta \quad (91)$$

where

$$\begin{aligned} \tilde{E}_\beta &\equiv \rho_\beta u_\beta^2 = n_\beta \bar{E}_\beta, \quad \tilde{T}_\beta \equiv \rho_\beta u_\beta^2 / 2 = n_\beta \bar{T}_\beta, \\ \tilde{U}_\beta &= \rho_\beta V_\beta'^2 = n_\beta \bar{U}_\beta = \rho_\beta u_\beta^2 / 2 \end{aligned} \quad (92)$$

One also obtains from (3)

$$\begin{aligned} \langle \mathbf{u}_\beta^2 \rangle &= \langle \mathbf{v}_\beta^2 \rangle + \langle \mathbf{V}_\beta'^2 \rangle + \langle 2\mathbf{v}_\beta \mathbf{V}_\beta' \rangle \\ &= \langle \mathbf{v}_\beta^2 \rangle + \langle \mathbf{V}_\beta'^2 \rangle \end{aligned} \quad (93)$$

since $\langle 2\mathbf{v}_\beta \mathbf{V}_\beta' \rangle = 0$, that can also be expressed as

$$\tilde{h}_\beta = \tilde{u}_\beta + p_\beta / \rho_\beta \quad (94)$$

and hence

$$H_\beta = U_\beta + p_\beta V_\beta \quad (95)$$

where V_β is the volume, and $H_\beta = M_\beta \tilde{h}_\beta$ and $U_\beta = M_\beta \tilde{u}_\beta$ are enthalpy and internal energy, respectively. Equation (95) represents the first law of thermodynamics

$$Q_\beta = U_\beta + W_\beta \quad (96)$$

when reversible heat and work are defined as $Q_\beta = H_\beta = T_\beta S_\beta$ and $W_\beta = p_\beta V_\beta$ at thermodynamic equilibrium and S_β is entropy [31].

For analysis of (87) one moves to the lower scale ($\beta-1$) and introduces the stretched internal time and space coordinates as

$$t_\beta = t_o + \varepsilon t_{\beta-1}, \quad z_\beta = z_o + \varepsilon z_{\beta-1} \quad (97)$$

where $\varepsilon \ll 1$ such that (87) becomes

$$\frac{\partial \Psi_\beta}{\partial t_{\beta-1}} + \mathbf{u}_\beta \nabla_{z_{\beta-1}} \Psi_\beta = 0 \quad (98)$$

By taking the first time derivative of (98) and substituting for $\partial \Psi_\beta / \partial t_{\beta-1}$ in the resulting equation from (98) itself and noting that $\mathbf{u}_\beta \neq f(t_{\beta-1}, z_{\beta-1})$ one arrives at the wave equation

$$\frac{\partial^2 \Psi_\beta}{\partial t_{\beta-1}^2} = u_\beta^2 \nabla_{z_{\beta-1}}^2 \Psi_\beta \quad (99)$$

or

$$\frac{\partial^2 \Psi_\beta}{\partial t^2} = u_\beta^2 \nabla^2 \Psi_\beta \quad (100)$$

where the subscripts ($\beta-1$) have been omitted. From substitution of the product solution $\Psi_\beta(\mathbf{x}, t) = \psi_\beta(\mathbf{x}) \Lambda_\beta(t)$ into (100) one obtains

$$\frac{\psi_\beta''}{\psi_\beta} = \frac{\Lambda_\beta''}{\Lambda_\beta u_\beta^2} = -\sigma_\beta^2 \quad (101)$$

where σ_β is the separation constant. The solution of the temporal part of (101) is

$$\Lambda_\beta = \exp(-i\sigma_\beta u_{o\beta} t) = \exp(-i\omega_\beta t) \quad (102)$$

suggesting that $\sigma_\beta u_\beta = \omega_\beta = 2\pi v_\beta$ is a circular frequency ω_β .

Following *Planck* [50, 51], one introduces by (24) the invariant expressions

$$\bar{E}_\beta \equiv m_\beta u_\beta^2 = h_\beta v_\beta, \quad \tilde{E}_\beta = n_\beta h_\beta v_\beta \quad (103)$$

when the harmonic atomic velocity is expressed as $u_\beta = \lambda_\beta v_\beta$. It was anticipated by *Dirac* [86] that amongst the fundamental constants (c , e , h), the *Planck* constant h might be expressible in terms of other fundamental constants (28). Using the result (103) one obtains

$$\sigma_\beta u_\beta = 2\pi \bar{E}_\beta / h_\beta \quad (104)$$

such that (102) becomes

$$\Lambda_\beta = \exp(-i2\pi \bar{E}_\beta / h_\beta) \quad (105)$$

By substitutions from (92), (104), and (32) into (101), one obtains the invariant time-independent *Schrödinger* equation [87, 88]

$$\nabla^2 \Psi_\beta + \frac{8\pi^2 m_\beta}{h^2} (\bar{E}_\beta - \bar{U}_\beta) \Psi_\beta = 0 \quad (106)$$

and through multiplication by (105), the invariant time-dependent *Schrödinger* equation

$$i\hbar_{\text{op}} \frac{\partial \Psi_\beta}{\partial t} + \frac{\hbar_\beta^2}{2m_\beta} \nabla^2 \Psi_\beta - \bar{U}_\beta \Psi_\beta = 0 \quad (107)$$

that governs the dynamics of particles from cosmic to tachyonic scales (Fig.1) [33, 88]. Since $\bar{E}_\beta = \bar{T}_\beta + \bar{U}_\beta$ by (92), the *Schrödinger* equation (106) gives the *stationary states* of particles that are trapped within *de Broglie* wave packet under the *potential* defined in (92) acting as *Poincaré* stress. In view of the definition (10), anticipation of an *external pressure* or *stress* as being the cause of particle stability by *Poincaré* [89-91] is a testimony to the true genius of this great mathematician, physicist, and philosopher.

It is emphasized that the stability of particle, *Heisenberg-Kramers* virtual oscillator [48], is due to the external force it experiences because of the pressure (92) induced by the peculiar velocity of the “atoms” of the field itself. In other words, it is the peculiar motions of atoms themselves that stabilizes different size atomic clusters. For example, peculiar motions of clusters are responsible for stability of eddies, and peculiar motions of molecules are responsible for stability of clusters, and so on. Hence, at the scale of stochastic electrodynamics $\beta = s$ ESD and chromodynamics $\beta = k$, EKD (see Figs. 1, 3) the model

suggests that peculiar motions of photons and tachyons are respectively responsible for the stability of electrons and photons. In view of the invariant *Planck* (44) and *Schrödinger* (107) equations, the model also suggests that *Planck* spectrum in equilibrium radiation represents stationary sizes of photon clusters, *de Broglie* molecules of light [92] or light bundles. Therefore, different “colors” of light are now recognized to correspond to different size photon clusters, photon wave packets, in accordance with the perceptions of *Newton* [53] and *Einstein* [66].

The formal derivation [33] of *Schrödinger* equation (107) from *Bernoulli* equation (76) provides for a new paradigm of the physical foundation of quantum mechanics. Soon after the introduction of his equation, *Schrödinger* himself [93] tried to identify some type of ensemble average interpretation of the wave function. Clearly, according to (85) the statistical ensemble nature of Ψ_β naturally arises from the velocity potential Φ'_β

Therefore, the dual and seemingly incompatible objective versus subjective natures of Ψ_β emphasized by *de Broglie* [1-3] can now be resolved. This is because the objective part of Ψ_β (85) is associated with the density ρ_β and accounts for the particle localization while the subjective part of Ψ_β is associated with the complex velocity potential Φ'_β that accounts for the observed action-at-a-distance as well as renormalization [94] and thereby the success of *Born's* [95] probabilistic interpretation of Ψ_β .

The definition of wave function (85) also helps to clarify the nature of wave-particle duality and *de Broglie's* theory of double solutions [1-3, 96, 97]. This is because the velocity potential Φ'_β acts as guidance wave that contains global information about the environment while remaining non-observable since it operates at the “hidden” “atomic” scale. Thus, the reason for success of separation of particle from wave according to *de Broglie* [1-3] and *Bohm and Vigier* [7, 9] becomes understood. Moreover, the reason for the failure of *von Neumann* no hidden variable theory [97-99] becomes clear since (107) for hierarchies of embedded statistical fields, Fig.1, could be subject to infinite hidden variables in harmony with *Gödel's* incompleteness theorem.

As one moves to smaller scales, one always finds a *continuum* because each element is composed of an entire statistical field (see Fig.1), such that one can again define a velocity potential $\Phi'_{\beta-1}$ and thereby define a new wave function $\Psi_{\beta-1} = -\rho_{\beta-1} \Phi'_{\beta-1}$. The cascade of particles as singularities embedded in

guidance waves is in exact agreement with the perceptions of *de Broglie* concerning interactions between the particle and the "hidden thermostat" [3]

"Here is another important point. I have shown in my previous publications that, in order to justify the well-established fact that the expression $|\psi(x,y,z,t)|^2 d\tau$ gives, at least with Schrödinger's equation, the probability for the presence of the particle in the element of volume $d\tau$ at the instant t , it is necessary that the particle jump continually from one guidance trajectory to another, as a result of continual perturbation coming from subquantal milieu. The guidance trajectories would really be followed only if the particle were not undergoing continual perturbations due to its random heat exchanges with the hidden thermostat. In other words, a Brownian motion is superposed on the guidance movement. A simple comparison will make this clearer. A granule placed on the surface of a liquid is caught by the general movement of the latter. If the granule is heavy enough not to feel the action of individual shocks received from the invisible molecules of the fluid, it will follow one of the hydrodynamic streamlines. If the granule is a particle, the assembly of the molecules of the fluid is comparable with the hidden thermostat of our theory, and the streamline described by the particle is its guiding trajectory. But if the granule is sufficiently light, its movement will be continually perturbed by the individual random impacts of the molecules of the fluid. Thus, the granule will have, besides its regular movement along one of the streamlines of the global flow of the fluid, a Brownian movement which will make it pass from one streamline to another. One can represent Brownian movement approximately by diffusion equation of the form $\partial\rho/\partial t = D\Delta\rho$, and it is interesting to seek, as various authors have done recently, the value of the coefficient D in the case of the Schrödinger equation corresponding to the Brownian movement.

I have recently studied ⁽¹⁴⁾ the same question starting from the idea that, even during the period of random perturbations, the internal phase of the particle remains equal to that of the wave. I have found the value $D = \hbar / (3m)$, which differs only by a numerical coefficient from the one found by other authors.

This concludes the account of my present ideas on the reinterpretation of wave mechanics with the help of images which guided me in my early work. My collaborators and I are working actively to develop these ideas in various directions. Today, I am convinced that the conceptions developed in the present article, when suitably developed and corrected at certain points, may in the future provide a real physical interpretation of present quantum mechanics."

To reveal the truly universal (Fig.1) nature of quantum mechanics one notes that for each statistical field shown in Figs.1 and 3 one can write a *Bernoulli* equation and by (76)-(107) arrive at a *Schrödinger* equation. At *Planck* scale $(\hbar G / c^3)^{1/2} \approx 10^{-35}$ m (see Fig.3) the hydrodynamic field represents the physical space [100] or *Casimir* vacuum [49] with its fluctuations and energy spectrum. According to definitions (27)-(28), the universal *Planck* and *Boltzmann* constants (h , k) respectively relate to spatial (λ) and temporal (ν) aspects of vacuum fluctuations. Also, at cosmic scales 10^{35} with $\beta = g$ (Figs.1, 3), the wave function Ψ_g will correspond to the wave function of the universe [101, 102]. The wave-particle duality of galaxies has been established by their observed quantized red shifts [103]. Indeed, it has been suggested by *Laughlin and Pines* [104] that *Schrödinger* equation may account for a part of the final theory, i.e. the theory of everything (TOE). Also, *Feynman et. al.* [105] suggested that *Schrödinger* equation may very possibly describe life itself.

In regards to the universality of quantum mechanics discussed above, it is interesting to examine the connections between classical and quantum mechanics. It seems that the new paradigm of quantum mechanics presented herein is in harmony with the perceptions of *Heisenberg* [106]

"We no longer say "Newtonian mechanics is false and must be replaced by quantum mechanics, which is correct." Instead we adopt the formula "Classical mechanics is a consistent self-enclosed scientific theory. It is strictly correct description of nature wherever its concepts can be applied"

This is because according to (76)-(107) quantum mechanics concerns behavior of virtual oscillators, wave packets in statistical fields such as in cosmology, hydrodynamics, molecular dynamics, electrodynamics or optics. Classical Newtonian mechanics is often concerned with motion of a few bodies such as the earth-moon-sun three-body problem.

According to Fig.3, a factor of 10^{-17} seems to separate the spatial scales of the stochastic fields of chromodynamics (10^{-35}), electrodynamics (10^{-17}), hydrodynamics (10^0), astrophysics (10^{17}), and cosmology (10^{35}). However, there are no mathematical or physical reasons to limit either large or small ends of the hierarchies shown in Fig.3. If one assumes with *Newton* that space is infinite and our universe is just one of many universes as suggested by Fig.3, then the recent observed asymmetry in the power spectrum from the right

versus the left side of our universe [107] schematically shown in Fig.20.




Fig.20 Asymmetry in measured cosmic power spectrum (a) calculated (b) measured [107].

appears to suggest that our universe is rotating in harmony with perceptions of *Kerr* [108] and *Gödel* [109] as well as all statistical fields of lower scales shown in Fig.1. Since it is known that galaxies including our Milky Way rotate, the cosmic iso-spin (12) is non-zero $\varpi_g \neq 0$. This suggests that the vorticity at cosmic scales (rotation of clusters of galaxies) may also be finite $\omega_g \neq 0$ that according to Fig.1 also means finite iso-spin of the many-universe field $\omega_g = \varpi_u \neq 0$. But finite vorticity of the multi-universe field $\omega_u \neq 0$ means rotation of our universe.

One may now introduce a new paradigm of the physical foundation of quantum mechanics according to which *Bohr's* stationary states will correspond to the statistically stationary sizes of clusters, *de Broglie* wave packets, which will be governed by *Maxwell-Boltzmann* distribution function (52) as shown in Fig.6 [33]. Next, different energy levels of quantum mechanics are identified as different size clusters (elements). For example, in EED field one views the transfer of a cluster from a small rapidly oscillating eddy (j) to a large slowly oscillating eddy (i) as transition from the high energy level (j) to the low energy level (i), see Fig.6, as schematically shown in Fig.21.




Fig.21 Transition of cluster c_{ij} from eddy-j to eddy-i leading to emission of molecule m_{ij} [34].

Such a transition will be accompanied with emission of a “sub-particle” that will be a

“molecule” (see Fig.21) to carry away the excess energy [34]

$$\Delta \varepsilon_{j\beta} = \varepsilon_{j\beta} - \varepsilon_{i\beta} = h(v_{j\beta} - v_{i\beta}) \quad (108)$$

in harmony with *Bohr's* theory of *atomic* spectra [48]. Therefore, the reason for the *quantum nature* of “molecular” energy spectra in equilibrium isotropic turbulence is that transitions can only occur between eddies with energy levels that satisfy the criterion of *stationarity* imposed by *Maxwell-Boltzmann* speed distribution function [100].

The new paradigm of quantum mechanics can help to resolve many of the central paradoxes of quantum mechanics. First is the problem of particle trajectory since according to the classical *Copenhagen* interpretation no trajectory could be defined for the particle during its transition that is absurd. According to the present model, any one of the particles of cluster (level) j could go through any trajectory to cluster (level) i and emit energy (108) with the exact same final consequence. The actual transition trajectories are dependent on the statistical nature of the distribution of clusters of various sizes (i.e energy levels). The probabilistic nature of various possible trajectories in effecting the same final outcome accounts for the success of *Feynman's* integral over all paths.

The next central problem of classical quantum mechanics is that of the collapse of the wave function [3]. Under the present model, the quantum mechanic wave function is related to the velocity potential (85), and hence to its conjugate stream function Ψ_β (65) that acts as *de Broglie* “guidance wave” [1-3, 96, 97] that guides the motion of the particle that is a singularity on the wave. Therefore, in the present paradigm the collapse of the wave function is identified as the collapse of the velocity potential due to the interactions between the measuring instrument and the flow field. In fact, the very dynamics of the wave function collapse could be systematically investigated and observed under various scenarios of experimental interactions with known velocity potentials.

Finally, the problems of double-slit experiment [2] as well as EPR [110] in classical quantum mechanics could now be resolved. This is because the EPR problem of *action at a distance* established by *Aspect et al.*, [111] could now be explained by possible superluminal tachyonic signals within physical space between the separated particles. Also, the double slit problem [2] is resolved due to the propagation of waves produced by the motion of particle in the tachyonic fluid that constitutes the

physical space. Simple experimental observations performed with macroscopic double-slit testing device under water employing small particles under micro-gravity is expected to verify the validity of the present model.

8 Compressibility of Physical Space and its impact on Special Theory of Relativity

The invariant time dependent *Schrödinger* equation (107) was derived from *Bernoulli* equation for an incompressible fluid. According to the scale-invariant statistical theory of fields schematically shown Fig.1 *physical space* is identified as a tachyonic fluid [100] that is *Dirac's* stochastic ether [112] or *de Broglie's* "hidden thermostat" [3]. Photons are considered to be composed of a large number of much smaller particles [100] called *tachyons* [113]. The importance of *Aristotle's* ether to the theory of electrons was emphasized by *Lorentz* [114, 115]

"I cannot but regard the ether, which is the seat of an electromagnetic field with its energy and its vibrations, as endowed with certain degree of substantiality, however different it may be from all ordinary matter"

Since the velocity of light is the mean thermal speed of tachyons, $u_k = c = v_t$, at least some of the tachyons must be *superluminal*.

The tachyonic fluid that constitutes the physical space is considered to be *compressible* in accordance with *Planck's* compressible ether [114]. If the compressible tachyonic fluid is viewed as an ideal gas, its change of density when brought isentropically to rest will be given by the expression involving *Michelson* number $Mi = v/c$ [116]

$$\rho = \rho_o \left[1 + \frac{\gamma - 1}{2} \frac{v^2}{c^2} \right]^{\frac{1}{\gamma - 1}} = \rho_o \left[1 + \frac{\gamma - 1}{2} Mi^2 \right]^{\frac{1}{\gamma - 1}} \quad (109)$$

With $\gamma = 4/3$ for photon gas, (109) leads to *Lorentz-FitzGerlad* contraction [116]

$$\lambda = \lambda_o \sqrt{1 - (v/c)^2} \quad (110)$$

that accounts for the null result of *Michelson-Morley* experiment [117].

Therefore, supersonic $Ma > 1$ (superchromatic $Mi > 1$) flow of air (tachyonic fluid) leads to the formation of *Mach* (*Poincaré-Minkowski*) cone that separates the zone of sound (light) from the

zone of silence (darkness). Compressibility of physical space can therefore result in *Lorentz-FitzGerald* contraction [114], thus accounting for relativistic effects [89-91, 114, 117-120] and providing a *causal explanation* [118] of such effects in accordance with the perceptions of *Poincaré* and *Lorentz* [89-91, 114, 120].

In view of the above considerations and in harmony with ideas of *Darrigol* [121] and *Galison* [122], one can identify two distinct paradigms of the Special Theory of Relativity [116]:

(A) Poincaré-Lorentz

Dynamic Theory of Relativity

Space and time (x, t) are altered due to causal effects of motion on the ether.

(B) Einstein

Kinematic Theory of Relativity

Space and time (x, t) are altered due to the two postulates of relativity:

- 1- The laws of physics do not change form for all inertial frames of reference.
- 2- Velocity of light is a universal constant independent of the motion of its source.

The result (110) and the almost constant cosmic temperature T_g and hence the speed of light

$$\lambda v = \lambda_o v_o = c(T_g) \approx c \quad (111)$$

lead to the frequency transformation

$$v = v_o / \sqrt{1 - (v/c)^2} \quad (112)$$

The relativistic transformation of frequency in (112) may also be expressed as contraction of time duration or transformation of period $\tau = 1/v$ as

$$\tau = \tau_o \sqrt{1 - (v/c)^2} \quad (113)$$

Hence, time durations and space extensions contract by (113) and (110) such that the speed of light remains invariant (111). It is emphasized however that the relation between time and space according to the dynamic theory of relativity of *Poincaré-Lorentz* is causal as emphasized by *Pauli* [118] and is induced by compressibility of physical space itself rather than being a purely kinematic effect as suggested by *Einstein* [119] according to paradigm (B) above.

Parallel to ideas of *Lorentz* [114], the concept of ether always played a crucial role in *Poincaré's* perceptions of relativity [121, 122] as he explicitly stated in his *Principle of Relativity* [123]

"We might imagine for example, that it is the ether which is modified when it is in relative motion in

“But the two times, past and future how can they be?, since the past is no more and the future is not yet?”

“Thus we can affirm that time is only in that it tends towards not-being”

“Yet Lord, we are aware of periods of time; we compare one period with another and say that some are longer, some shorter”

“Does not my soul speak truly to You when I say that I can measure time? For so it is, O Lord my God. I measure it and I do not know what it is that I am measuring”

Now, the classical problem of how could a finite time be constructed from multitudes of instants, “nows”, that do not exist can be addressed by the invariant definition of atomic time in (114c)

$$\tau_{\beta} = \sum t_{\beta} = \sum \tau_{\beta-1} \quad (116)$$

This is because the “atomic” instant $t_{\beta} = 0$ of scale β has a finite duration $\tau_{\beta-1}$ at the lower scale $\beta-1$ (114c). Thus, one uses clocks of $\beta-1$ scale to measure time of β scale, clocks of $\beta-2$ scale to measure time of $\beta-1$ scale, and so on at infinitum. Therefore, similar considerations employed for description of analysis in space continuum [127] will be required to describe the temporal continuum discussed by Weyl [128]

“Exact time- or space-points are not the ultimate, underlying, atomic elements of the duration or extension given to us in experience. On the contrary, only reason, which thoroughly penetrates what is experientially given, is able to grasp exact ideas”

Accordingly, if one does not wish to allow for infinite divisibility of time and space, then following *Leibniz* [127] one must introduce the *temporal monad* just like the *spatial monad* to represent the absolute smallest “atom” of time or *Chronon*. Also, parallel to *Heisenberg’s spatial uncertainty principle* [129]

$$\Delta \lambda_{\beta} \Delta p_{\beta} \geq h \quad (117)$$

that limits the resolution of spatial measurements, the *temporal uncertainty principle* [100]

$$\Delta \nu_{\beta} \Delta p_{\beta} \geq k \quad (118)$$

limits the resolution of time measurements.

Since time is identified as a physical attribute of the dynamics of tachyonic atoms (114c) and space is identified as this compressible tachyonic fluid itself, it is clear that the causal connections between space and time in relativistic physics become apparent. For example, in the classical problem of *twin paradox* of the special theory of

relativity, the different times experienced by the twins could be attributed to the different rates of biological reactions in their body induced by the compressibility of physical space. According to the causal dynamic theory of relativity of *Poincaré-Lorentz*, the reason for the coincidence of directions of biological and cosmological times [130] becomes apparent.

The physical space or *Casimir* vacuum [49] when identified as a compressible tachyonic fluid provides a new paradigm for the physical foundation of quantum gravity [100]. The general implications of the model to time reversibility in quantum cosmology and to *Everett’s* many-universe theory [101, 131-132] require further future investigations. Because of the definition of atomic time in (114c), quantum theories of gravity [131-138] may have wave functions Ψ_g that instead of *Wheeler-DeWitt* equation [101, 131, 132, 133, 138]

$$H\Psi_g = 0 \quad (119)$$

satisfy the *modified Wheeler-DeWitt* equation

$$i\hbar \frac{\partial \Psi_g}{\partial t_{\beta-1}} = H\Psi_g \quad (120)$$

that is *Schrödinger* equation (107). The resurrection of time in (120) is made possible because the new “atomic” time arises from internal degrees of freedom, permitting $\Psi_g(x_1, x_2, x_3, t_{\beta}, t_{\beta-1})$ and the associated $g_{ij}(x_1, x_2, x_3, t_{\beta}, t_{\beta-1})$, that by thermodynamic considerations is related to the temperature of the field [100].

The definition of time in (114) is in accordance with *ephemeris* time in astronomy [136, 139]

$$\delta t = \sqrt{\frac{\sum_i m_i (\delta d_i)^2}{2(E-V)}} \quad (121)$$

Substituting in (121) the kinetic energy $K = E - V = mv^2 / 2$ and mass fraction $Y_i = m_i / m$ results in

$$\begin{aligned} \delta t &= \sqrt{\frac{\sum_i m_i (\delta d_i)^2}{2(E-V)}} = \sqrt{\frac{m \sum_i Y_i (\delta d_i)^2}{mv^2}} \\ &= \sqrt{\frac{m(\delta d)^2}{mv^2}} = \frac{\delta d}{v} \end{aligned} \quad (122)$$

that is in accordance with (114). The mean extension δd and duration δt are mass-average of the component extension δd_i and duration δt_i and the corresponding velocity defined as

$$\delta t_i = \sqrt{\frac{m_i (\delta d_i)^2}{2(E_i - V_i)}} \quad , \quad v_i = \frac{\delta d_i}{\delta t_i} \quad (123)$$

where $E_i = K_i + V_i$, $K_i = m_i v_i^2 / 2$, and $V_i = -Gm_i m_j / r_{ij}$. The invariant definition of time in (114) suggests that the description of temporal continuum [128] requires introduction of “temporal measure” that relates to the important problems of duration and simultaneity identified by *Poincaré* in 1898 [140] as emphasized by *Barbour* [136].

9 Implications to Dirac Relativistic Wave Equation

It is interesting to explore the implications of the quantum mechanical wave function (85) to *Dirac's* relativistic wave equation [141]. To do this one notes that since the peculiar velocity of particle at scale β

$$\mathbf{V}'_\beta = \mathbf{u}_\beta - \mathbf{v}_\beta \quad , \quad (124)$$

is the diffusion velocity of scale $(\beta-1)$ one can express (124) in terms of the velocities of the lower scale $\beta-1$ as in (3)

$$\mathbf{V}_{\beta-1} = \mathbf{v}_{\beta-1} - \mathbf{w}_{\beta-1} \quad (125)$$

Thus, the results (124) and (85) relate the quantum mechanic wave function to the equation of motion at scale $(\beta-1)$ through diffusion velocity (125).

Hence, one starts with *Cauchy* equation of motion (7) at scale $(\beta-1)$

$$\frac{\partial \mathbf{p}_{i\beta-1}}{\partial t} + \frac{\partial}{\partial x_j} (\mathbf{p}_{i\beta-1} \mathbf{v}_{j\beta-1}) = -\nabla \cdot \mathbf{P}_{ij\beta-1} \quad (126)$$

Next, the pressure within the particle at the lower scale $(\beta-1)$ is considered to be constant such that $\nabla \cdot \mathbf{P}_{ij\beta-1} = 0$ and (126) reduces to

$$\frac{\partial \mathbf{p}_{i\beta-1}}{\partial t} + \frac{\partial}{\partial x_j} (\mathbf{p}_{i\beta-1} \frac{\partial x_j}{\partial t}) = 0 \quad (127)$$

Parallel to derivation of *Schrödinger* equation [33], one next considers the moving coordinates

$$\mathbf{x}_{j\beta} = \mathbf{z}_{j\beta} + \alpha_j \mathbf{w}_{\beta} t \quad (128)$$

that for uniform velocity \mathbf{w}_β results in

$$\mathbf{v}_{j\beta-1} = \mathbf{V}_{j\beta-1} + \alpha_j \mathbf{w}_{\beta-1} = \mathbf{V}_{j\beta-1} + \mathbf{w}_{j\beta-1} \quad (129)$$

in accordance with (125). The components of convective velocity in (128) are expressed as

$$\mathbf{w}_{j\beta-1} = \alpha_j \mathbf{w}_{\beta-1} \quad (130)$$

such that

$$\mathbf{w}_{\beta-1}^2 = \sum \alpha_j^2 \mathbf{w}_{\beta-1}^2 = \sum \mathbf{w}_{j\beta-1}^2 \quad (131)$$

For uniform convection velocity (130) by (125) and substitution from (13c) one obtains from (127)

$$\frac{\partial \mathbf{p}_{i\beta-1}}{\partial t} + \alpha_j \mathbf{w}_{\beta-1} \frac{\partial \mathbf{p}_{i\beta-1}}{\partial x_j} = v_\beta \frac{\partial^2 \mathbf{p}_{i\beta-1}}{\partial x_j^2} \quad (132)$$

that is identical to (19) when $\nabla \mathbf{p}_{\beta-1} = \nabla \cdot \mathbf{v}_{\beta-1} = 0$.

For uniform convective velocity \mathbf{W}_β (125) gives

$$\mathbf{v}_{i\beta-1} = \mathbf{w}_{i\beta-1} + \mathbf{V}_{ij\beta-1} = A + \mathbf{V}'_{ij\beta} \quad (133)$$

where A is a constant. Also, for an incompressible fluid with $\rho_{\beta-1} A = B$ one obtains from (133)

$$\rho_{\beta-1} \mathbf{v}_{i\beta-1} = B + \rho_{\beta-1} \mathbf{V}'_{ij\beta} = B + Y_{\beta-1} \nabla_i \rho_\beta \Phi'_\beta \quad (134)$$

that by (85) gives

$$\mathbf{p}_{i\beta-1} = B + Y_{\beta-1} \nabla_i \Psi_\beta \quad (135)$$

Substituting from (135) into (132) leads to

$$\frac{\partial \Psi_\beta}{\partial t} + \alpha_j \mathbf{w}_{\beta-1} \frac{\partial \Psi_\beta}{\partial x_j} = \alpha_m \frac{\partial^2 \Psi_\beta}{\partial x_j^2} \quad (136)$$

where the viscosity tensor α_m by (13c) is defined as

$$v_\beta = v_{ij\beta} = \alpha_m \quad , \quad \mathbf{V}_{ij\beta} = -v_{ij\beta} \frac{\partial \ln \mathbf{p}_{j\beta}}{\partial x_i} \quad (137)$$

Following *Dirac* [141], by (137), and (125)-(131) the parameters (α_j, α_m) are considered to be tensors representing 4×4 matrices. However, since according to *Dirac* one of his four equations is redundant [141], the final wave equation will only involve three components such that by (131)

$$\alpha_1^2 = \alpha_2^2 = \alpha_3^2 = 1 \quad (138)$$

in accordance with *Dirac* theory [141]. Also, the definitions (131) and (137) are in harmony with the perceptions of *Dirac* [141] who anticipated that (α_j, α_m) may be related to some coordinates associated with internal degrees of freedom.

Relating *Schrödinger* and *Dirac* wave functions by the expression $\Psi_\beta = \tilde{\Psi}_\beta(z_j) \Psi_{D\beta}(x_j, t)$ with

$$\tilde{\Psi}_\beta(z_j) = \exp[(\sqrt{im_{\beta-1} \mathbf{w}_{\beta-1}^2 / \hbar}) z_j] \quad (139)$$

, $\Psi_{D\beta} = \exp(i\alpha_m E_\beta t / \hbar) \exp(-i\alpha_m \alpha_j \mathbf{w}_{\beta-1} x_j / \hbar)$, and the total energy defined as $E_{\beta-1} = 2m_{\beta-1} \mathbf{w}_{\beta-1}^2$, one obtains

from (128), (139) and (136) the scale invariant relativistic wave equation

$$i\hbar\left(\frac{1}{w_{\beta-1}}\frac{\partial\Psi_{D\beta}}{\partial t}+\alpha_j\frac{\partial\Psi_{D\beta}}{\partial x_j}\right)+(\alpha_m m_{\beta-1}w_{\beta-1})\Psi_{D\beta}=0 \quad (140)$$

At the important scale of *Casimir* vacuum [49] (ETD in Fig.1), when $w_{\beta-1} = v_t = u_k = c$ is the speed of light, equation (140) becomes *Dirac* relativistic wave equation for electrons [141]

$$i\hbar\left(\frac{1}{c}\frac{\partial\Psi_{D\beta}}{\partial t}+\alpha_j\frac{\partial\Psi_{D\beta}}{\partial x_j}\right)+(\alpha_m mc)\Psi_{D\beta}=0 \quad (141)$$

Because it was by sheer genius and mathematical intuition that *Dirac* arrived at his relativistic wave equation (141), the physical basis of this equation remains quite abstract and mysterious. Therefore, the simple derivation of *Dirac* equation presented above may help the understanding of this important equation. It is also noted that the wave function Ψ_β in (85) remains well defined even in the presence of spin as long as $\nabla \times \mathbf{v}_\beta = \nabla \times \mathbf{u}_\beta$ such that $\nabla \times \mathbf{V}'_\beta = 0$ by (124). The true significance of the tensors (α_j, α_m) in (130) and (137) as well as the wave function (139) require further future investigations.

10 Concluding Remarks

A modified statistical theory of turbulence was presented and the connections between the problems of turbulence and quantum mechanics were further explored. New paradigms for physical foundations of invariant *Planck* law, *Schrödinger* equation, and *Dirac* relativistic wave equation were presented. The predicted velocity profiles for flow over a flat plate were compared with measurements for LED, LCD, LMD, and LAD scales. The universal nature of turbulence across broad range of spatio-temporal scales is in harmony with occurrence of fractals in physical science emphasized by *Takayasu* [142].

Acknowledgments: This research was in part supported by NASA grant No. NAG3-1863.

References:

- [1] Broglie, L. de, *C. R. Acad. Sci., Paris*, **183**, 447 (1926); **184**, 273 (1927); **185**, 380 (1927); *Nature* **118** (2969) 441 (1926).
- [2] Broglie, L. de, *Non-Linear Wave Mechanics, A Causal Interpretation*, Elsevier, New York, 1960.
- [3] Broglie, L. de, *Found. Phys.* **1**, 5 (1970).
- [4] Madelung, E., *Z. Physik*, **40**, 332 (1926).
- [5] Schrödinger, E., *Berliner Sitzungsberichte*, **144** (1931).
- [6] Fürth, R., *Z. Phys.* **81**, 143 (1933).
- [7] Bohm, D., *Phys. Rev.* **85**, 166 (1952).
- [8] Takabayasi, T., *Prog. Theor. Phys.* **70**, 1 (1952).
- [9] Bohm, D., and Vigier, J. P., *Phys. Rev.* **96**, 208 (1954).
- [10] Nelson, E. *Phys. Rev.* **150**, 1079 (1966).
- [11] Nelson, E. *Quantum Fluctuations*, Princeton University Press, Princeton, New Jersey, 1985.
- [12] de la Peña, L., *J. Math. Phys.* **10**, 1620 (1969).
- [13] de la Peña, L., and Cetto, A. M., *Found. Phys.* **12**, 1017 (1982).
- [14] Barut, A. O., *Ann. Physik*, **7**, 31 (1988).
- [15] Barut, A. O., and Bracken, A. J., *Phys. Rev. D* **23**, 2454 (1981).
- [16] Vigier, J. P., *Lett. Nuvo Cim.* **29**, 467 (1980); Gueret, Ph., and Vigier, J. P., *Found. Phys.* **12**, 1057 (1982); Cufaro Petroni, C., and Vigier, J. P., *Found. Phys.* **13**, 253 (1983); Vigier, J. P., *Found. Phys.* **25**, 1461 (1995).
- [17] Arecchi, F. T., and Harrison, R. G., *Instabilities and Chaos in Quantum Optics*, Springer-Verlag, Berlin 1987.
- [18] Reynolds, O., *Phil. Trans. Roy. Soc. A* **186**, 123, (1895).
- [19] Taylor, G. I., I-IV, *Proc. Roy. Soc. A* **151**, 421 (1935).
- [20] Kármán, T. von, and Howarth, L., *Proc. Roy. Soc. A* **164**, 192 (1938).
- [21] Robertson, H. P., *Proc. Camb. Phil. Soc.* **36**, 209 (1940).
- [22] Kolmogoroff, A. N., *C. R. Acad. Sci. U. R. S. S.* **30**, 301 (1941); **32**, 16 (1942). *J. Fluid Mech.* **13**, 82 (1962).
- [23] Obukhov, A. M., *C. R. Acad. Sci. U. R. S. S.* **32**, 19 (1941); *J. Fluid Mech.* **13**, 77 (1962).
- [24] Chandrasekhar, S., *Rev. Mod. Phys.* **15**, 1 (1943).
- [25] Chandrasekhar, S., *Stochastic, Statistical, and Hydrodynamic Problems in Physics and Astronomy*, Selected Papers, vol.3, University of Chicago Press, Chicago, 1989.
- [26] Heisenberg, W., *Proc. Roy. Soc. A* **195**, 402 (1948); *Z. Phys.* **124**, 628 (1948).
- [27] Batchelor, G. K., *The Theory of Homogeneous Turbulence*, Cambridge University Press, Cambridge, 1953.
- [28] Landau, L. D., and Lifshitz, E. M., *Fluid Dynamics*, Pergamon Press, New York, 1959.
- [29] Tennekes, H., and Lumley, J. L., *A First Course In Turbulence*, MIT Press, 1972.
- [30] Sohrab, S. H., *Proceeding of the 31st ASME National Heat Transfer Conference*, HTD-Vol. **328**, 37-60 (1996).
- [31] Sohrab, S. H., *Rev. Gén. Therm.* **38**, 845 (1999).
- [32] Sohrab, S. H., In: *Theoretical and Experimental Aspects of Fluid Mechanics*, S. H. Sohrab, H. C. Catrakis, and F. K. Benra (Eds.), WSEAS Press, 27-35, 2008.

- [33] Sohrab, S. H., *WSEAS Transactions on Mathematics* **6** (2), 254 (2007).
- [34] Sohrab, S. H., In: *Recent Advances in Fluid Mechanics & Aerodynamics*, S. Sohrab, H. Catrakis, and N. Kobasko (Eds.), pp:134-140, WSEAS Press, 2009, ISBN: 978-960-474-106-9.
- [35] de Groot, R. S., and Mazur, P., *Nonequilibrium Thermodynamics*, North-Holland, 1962.
- [36] Schlichting, H., *Boundary-Layer Theory*, McGraw Hill, New York, 1968.
- [37] Williams, F. A., *Combustion Theory*, Benjamin Cummings, New York, 1985.
- [38] Chapman, S., and Cowling, T. G., *The Mathematical Theory of Non-uniform Gases*, Cambridge University Press, Cambridge, 1953.
- [39] Hirschfelder, J. O., Curtiss, C. F., and Bird, R. B., *Molecular Theory of Gases and Liquids*, Wiley, New York, 1954.
- [40] Sohrab, S. H., In: *New Aspects of Fluid Mechanics and Aerodynamics*, S. H. Sohrab, H. J. Catrakis, N. Kobasko, N., Necasova, and N. Markatos (eds.) WSEAS press, 2008.
- [41] Sohrab, S. H., O., In: *Recent Advances in Systems*, N. Mastorakis, V. Mladenov, Z. Bojkovic, S. Kartalopoulos, A. Varonides, and M. Jha (Eds.), pp: 557-568, WSEAS Press, 2009, ISBN: 978-960-474-097-0.
- [42] Lugt, H. J., *Vortex Flow in Nature and Technology*, Wiley, New York, 1983.
- [43] Bahatia, A. B., and Singh, R. N., *Mechanics of Deformable Media*, IOP publication, Boston, 1986.
- [44] Carrier, G. On slow viscous flow, *Nonr-653-00-1*, (1953).
- [45] Olmstead, W. E. Reciprocity relationships in viscous hydrodynamics, *Acta Mechanica*. **21**, 289 (1975).
- [46] Roper, M., and Brenner, M. P., *PNAS* **106**, 2977 (2009).
- [47] Townsend, A. A., *Proc. Camb. Phil. Soc.* **43**, 560 (1947).
- [48] van der Waerden, B. L., *Sources of Quantum Mechanics*, B. L. van der Waerden (ed.), Dover, New York, 1967.
- [49] Casimir, H. B. G., *Proc. K. Ned. Akad. Wet.* **51**, 793 (1948).
- [50] Planck, M., *The Theory of Heat Radiation*, Dover, New York, 1991.
- [51] Planck, M., *Ann. der Physik* **4**, 553 (1901).
- [52] Jackson, J. D., *Classical Electrodynamics*, Wiley, New York, 1975.
- [53] Newton, I., *Optics*, Dover, New York, 1952.
- [54] Broglie, L., *Le Mechanique Ondulatoire du photon* **1**, Paris: Herman et Cie, 1940.
- [55] Bass, L., and Schrödinger, E., *Proc. Roy. Soc. A* **232**, 1-6 (1955).
- [56] Vigier, J. P., *Found. Phys.* **24**, 61 (1994).
- [57] de Beauregard, O. Costa., In: *Waves and Particles in Light and Matter*. Alwyn van der Merwe and Augusto Garuccio (Eds.), Plenum Press, New York, 1994.
- [58] Evans, M. W., In: *Waves and Particles in Light and Matter*. Alwyn van der Merwe and Augusto Garuccio (Eds.), p.259, Plenum Press, New York, 1994.
- [59] Sohrab, S. H., O., in *New Aspects of Heat Transfer, Thermal Engineering, and Environment*, S. H. Sohrab, H. J. Catrakis, N. Kobasko, (Eds.), pp: 434-442, WSEAS Press, 2008, ISBN: 978-960-6766-97-8.
- [60] Sohrab, S. H., *Bull. Am. Phys. Soc.* **49** (1) 255 (2004).
- [61] Chandrasekhar, S., *Newton's Principia for the Common Reader*, Oxford University Press, New York, 1995.
- [62] Pauli, W., *Pauli Lectures on Physics*, Vol.3, p.14, MIT Press, 1973.
- [63] Long, C. A., and Sohrab, S. H., In: *Recent Advances on Applied Mathematics*, Long, C. A., Sohrab, S. H., Bogнар, G., and Perlovsky, L., (Eds.), p.87-97, WSEAS Press, 2008.
- [64] De Pretto, O., Ipotesi dell'etere nella vita dell'universo. *Atti del Reale Inst. Veneto di Scienze, Lettere ed Arti.* **63** (2), 439-499 (1904).
- [65] Rayleigh, Lord., *Phil. Mag.*, **49**, 539 (1900).
- [66] Einstein, A., *Ann. Phys. (Leipzig)* **17**, 132 (1905).
- [67] Sonntag, R. E., Van Wylen, G. E., *Fundamentals of Statistical Thermodynamics*, Wiley, New York, 1966.
- [68] Van Atta, C. W., and Chen, W. Y., *J. Fluid Mech.* **38**, 743 (1969).
- [69] Landahl, M. T., and Mollo-Christensen, E., *Turbulence and Random Processes in Fluid Mechanics*, Cambridge University Press, Cambridge, 1992.
- [70] Darrigol, O., *World of Flow*, Oxford University Press, New York, 2005.
- [71] Lumley, L. J., Berkooz, G., Elezgaray, J., Holmes, P., Poje, A., and Volte, C., In: *Simulation and Modeling of Turbulent Flows*. Gatski, B. T., Hussaini, M. Y., Lumley, L. J. (Eds), Oxford University Press 1996.
- [72] McComb, W. D., and Shanmugasundaram, V., *J. Fluid Mech.* **143**, 95 (1984).
- [73] Saddoughi, G. S., and Veeravalli, V. S., *J. Fluid Mech.* **268**, 333 (1994).
- [74] Marusic, I., Mathis, R., and Hutchins, N., *Science* **329**, 193 (2010).
- [75] Lin, C. C., *Turbulent Flows and Heat Transfer*, Princeton University Press, Princeton, 1959.
- [76] Ling, S. C., and Huang, T. T., *J. Phys. Fluids.* **13**, 2912 (1970).
- [77] Chandrasekhar, S., Turbulence- A physical theory of astrophysical interest, in [25], (1949).
- [78] Martinelli, R. C., *Transactions, ASME* **69**, 947 (1947).
- [79] Ueda, H., and Hinze, O. J., *J. Fluid Mech.* **67**, 125 (1975).

- [80] Zagarola, M. V., Perry, A. E., and Smits, A. J., Log laws or power laws: The scaling in the overlap region, *Phys. Fluids* **9** (7), 2094 (1997).
- [81] Lancien, P., Lajeunesse, E., and Metivier, F., Near wall velocity measurements by particle-tracking, arXiv:physics/0701131v1 [physics.flu-dyn], 11 January 2007.
- [82] Meinhardt, C. D., Wereley, S. T., and Santiago, J. G., *Experiments in Fluids* **27**, 414 (1999).
- [83] Sommerfeld, A., *Mechanics of Deformable Bodies*, Academic Press, New York, 1950.
- [84] Bognar, G., and Sohrab, S. H., In: *Recent Advances in Fluid Mechanics & Aerodynamics*, S. Sohrab, H. Catrakis, and N. Kobasko (Eds.), pp: 41-48, WSEAS Press, 2009, ISBN: 978-960-474-106-9.
- [85] Sohrab, S. H., *Bull. Am. Phys. Soc.* **43** (1) 781 (1998).
- [86] Dirac, P. A. M., *Sci. American* **208**, 45 (1963).
- [87] Schrödinger, E., *Ann. Physik* **79**, 361 (1926); **81**, 109 (1926).
- [88] Sohrab, S. H., *WSEAS Transactions on Mathematics* **4**, Vol 3, 764 (2004); *WSEAS Transactions on Circuits and Systems* **4**, Vol 3, 1017 (2004).
- [89] Poincaré, H., Sur la dynamique de l'électron, *C. R. Acad. Sci. Paris* **140**, 1505 (1905).
- [90] Poincaré, H., Sur la dynamique de l'électron, *Rend. Circ. Mat. Palermo* **21**, 129 (1906).
- [91] Logunov, A. A., *On the Articles by Henri Poincaré, "On the Dynamics of the Electron"*, Dubna: JINR, 2001.
- [92] Broglie, L. de, *Matter and Light-The New Physics*, Dover, New York, 1937.
- [93] Schrödinger, E., *Phys. Rev.* **28**, 1049 (1926).
- [94] 't Hooft, G., *Rev. Mod. Phys.* **72** (2), 333 (1999).
- [95] Born, M., *Z. Physik* **37**, 863 (1926).
- [96] Vigier, J. P., Louis de Broglie—Physicist and Thinker, In: *Quantum, Space and Time-The Quest Continues*, Asim O. Barut, Alwyn van der Merwe, and Jean-Pierre Vigier (Eds.), Cambridge University Press, 1984.
- [97] Lochak, G., In: *Quantum, Space and Time-The Quest Continues*, Asim O. Barut, Alwyn van der Merwe, and Jean-Pierre Vigier (Eds.), pp.11-33, Cambridge University Press, 1984.
- [98] Polkinghorne, J. C., *The Quantum World*, Princeton University Press, 1984.
- [99] Bell, J. S., In: *Quantum, Space and Time-The Quest Continues*, Asim O. Barut, Alwyn van der Merwe, and Jean-Pierre Vigier (Eds.), pp.66-76, Cambridge University Press, 1984.
- [100] Sohrab, S. H., *WSEAS Transactions on Mathematics* **4**, Vol.3, 764 (2004).
- [101] DeWitt, S. B., *Phys. Rev.* **160**, 1113 (1967).
- [102] Hartle, J. B., and Hawking, S. W., *Phys. Rev. D* **28**, 2960 (1983).
- [103] Dersarkissian, M., *Lett. Nuovo Cim.* **40**, 390 (1984).
- [104] Laughlin, R. B., and Pines, D., *PNAS* **97** (1) 28-31 (2000).
- [105] Feynman, R., Leighton, R. B., and Sands, M., *Lectures on Physics*, Vol. II, p.41-11, Addison-Wesley Publishing, New York, 1964.
- [106] Bokulich, A., *Reexamining the Quantum-Classical Relation*, Cambridge University Press, New York, 2008.
- [107] Kamionkowski, M., Is the universe (statistically) isotropic? Lecture delivered at physics department, Northwestern University, Evanston, Illinois, March (2010).
- [108] Kerr, R. P., *Phys. Rev. Let.* **11** (5), 237 (1963).
- [109] Gödel, K., *Rev. Mod. Phys.* **21** (3), 447 (1949).
- [110] Einstein, A., Podolsky, B., and Rosen, N., *Phys. Rev.* **41**, 777 (1935).
- [111] Aspect, A., Dalibard, J., Gerard, R., *Phys. Rev. Let.* **49**, (25) 1804 (1982).
- [112] Dirac, P. A. M., *Nature* **168**, 906 (1951).
- [113] Recami, E., *Riv. Nuovo Cim.* **9**, No.6, (1986); *Tachyons, Monopoles and Related Topics*, North-Holland, Amsterdam, 1978.
- [114] Lorentz, H. A., *Selected Works of L.H. Lorentz*, vol.5, Nersessian, N. J., and Cohen, H. F., (Eds), Palm Publications, Nieuwerkerk, 1987, p.7.
- [115] Kox, A. J., in *Einstein and the History of General Relativity*, Howard, D., and Stachel, J. (Eds.), pp.201-212, Birkhäuser, Boston, 1989.
- [116] Sohrab, S. H., in *New Aspects of Heat Transfer, Thermal Engineering, and Environment*, S. H. Sohrab, H. J. Catrakis, N. Kobasko, (Eds.), pp: 434-442, WSEAS Press, 2008, ISBN: 978-960-6766-97-8.
- [117] Michelson A. A., and Morley, E. W., On the relative motion of the earth and luminiferous ether, *Amer. J. Sci.* **34**, 333-345 (1887).
- [118] Pauli, W., *Theory of Relativity*, Dover, New York, 1958.
- [119] Einstein, A., *Ann. Phys. (Leipzig)* **18**, 639 (1905).
- [120] Whittaker, E. T., *A History of the Theories of Aether and Electricity*, vol.2, Tomash Publishers, New York, 1987.
- [121] Darrigol, O., *Isis* **95**, 614-626 (2004).
- [122] Galison, P., *Einstein's Clocks, Poincaré's Maps*, W. W. Norton & Company, New York, 2003.
- [123] Poincaré, H., *Bulletin des Sciences Mathématiques* **28**, 302-324 (1904), English translation in: *The Monist* XV, No.1, 1 (1905).
- [124] Poincaré, H., *Arch. Neerland.* **5**, 252 (1900). <http://www.physicsinsights.org/poincare-1900.pdf>.
- [125] Poincaré, H., *Mathematics and Science Last Essay*, p.19, General Books, Memphis, Tennessee, 2010.
- [126] Aristotle, In: *Time*, Westphal J. and Levenson, C., (Eds.), Hackett Publishing Company, Indianapolis, Indiana, 1993.
- [127] Bell, L. J., *The Continuous and the Infinitesimal in Mathematics and Philosophy*, Polimetria, Milano, Italy, 2006.
- [128] Weyl, H., *The Continuum*, The Thomas Jefferson University Press, Kirksville, MO, 1987.

- [129] Heisenberg, W., *The Physical Principles of Quantum Theory*, Dover, New York, 1949.
- [130] Hawking, S. W., *A Brief History of Time*, Bantam Books, 1988; *Phys. Rev. D* **32** (10), 2489 (1985).
- [131] DeWitt, S. B., *Physics Today*, September, (1970).
- [132] Wheeler, J. A., Superspace and the nature of quantum geometro-dynamics. In: *Battelle Rencontres*, edited by C. M. DeWitt and J. A. Wheeler, 242-307, Benjamin, New York 1968.
- [133] Ashtekar, A., *Current Science* **89** (12), 2064 (2005).
- [134] Kiefer, C., Does Time Exist in Quantum Gravity? www.fqxi.org/communitiy/forum/topic/265 (2008).
- [135] Rovelli, C., Forget Time, www.fqxi.org/communitiy/forum/topic/237 (2008).
- [136] Barbour, J., The Nature of Time, www.fqxi.org/communitiy/forum/topic/360 (2008).
- [137] Smolin, L., *The Life of the Cosmos*, Oxford University Press, Oxford, 1997; In: *Conceptual Problems of Quantum Gravity*, Ashtekar, A., and Stachel, J. (eds.), Birkhäuser, Boston, 1991.
- [138] Zeh, H. D., *The Physical Basis of the Direction of Time*, Springer-Verlag, 1999.
- [139] Clemence, G. M., *Rev. Mod. Phys.* **29**, 2 (1957).
- [140] Poincaré, H., *Rev. Metaphys. Morale* **6**, 1 (1898); English translation; "The measure of time," in: Poincaré, H., *The Value of Science*.
- [141] Dirac, A. M. P., *Directions in Physics*, Wiley, New York, 1978.
- [142] Takayasu, H., *Fractals in Physical Science*, Wiley, New York, 1990.